

A variational formulation for frame-based inverse problems

Caroline Chaux¹, Patrick L Combettes², Jean-Christophe Pesquet¹
and Valérie R Wajs²

¹ Institut Gaspard Monge and UMR CNRS 8049, Université de Marne la Vallée,
77454 Marne la Vallée Cedex 2, France

² Laboratoire Jacques-Louis Lions—UMR CNRS 7598, Faculté de Mathématiques,
Université Pierre et Marie Curie—Paris 6, 75005 Paris, France

E-mail: plc@math.jussieu.fr

Received 15 December 2006, in final form 15 May 2007

Published 14 June 2007

Online at stacks.iop.org/IP/23/1495

Abstract

A convex variational framework is proposed for solving inverse problems in Hilbert spaces with *a priori* information on the representation of the target solution in a frame. The objective function to be minimized consists of a separable term penalizing each frame coefficient individually, and a smooth term modelling the data formation model as well as other constraints. Sparsity-constrained and Bayesian formulations are examined as special cases. A splitting algorithm is presented to solve this problem and its convergence is established in infinite-dimensional spaces under mild conditions on the penalization functions, which need not be differentiable. Numerical simulations demonstrate applications to frame-based image restoration.

1. Introduction

In inverse problems, certain physical properties of the target solution \bar{x} are most suitably expressed in terms of the coefficients $(\bar{\xi}_k)_{k \in \mathbb{K} \subset \mathbb{N}}$ of its representation $\bar{x} = \sum_{k \in \mathbb{K}} \bar{\xi}_k e_k$ with respect to a family of vectors $(e_k)_{k \in \mathbb{K}}$ in a Hilbert space $(\mathcal{H}, \|\cdot\|)$. Traditionally, such linear representations have been mostly centred on orthonormal bases such as, for instance, in Fourier, wavelet or bandlet decompositions [8, 25, 26]. Recently, attention has shifted towards more general, overcomplete representations known as *frames* (see [6, 7, 16, 20, 33] for specific examples). Recall that a family of vectors $(e_k)_{k \in \mathbb{K}}$ in \mathcal{H} constitutes a frame if there exist two constants μ and ν in $]0, +\infty[$ such that

$$(\forall x \in \mathcal{H}) \quad \mu \|x\|^2 \leq \sum_{k \in \mathbb{K}} |\langle x | e_k \rangle|^2 \leq \nu \|x\|^2. \quad (1.1)$$

The associated frame operator is the injective bounded linear operator

$$F : \mathcal{H} \rightarrow \ell^2(\mathbb{K}) : x \mapsto ((x | e_k))_{k \in \mathbb{K}}, \quad (1.2)$$

the adjoint of which is the surjective bounded linear operator

$$F^* : \ell^2(\mathbb{K}) \rightarrow \mathcal{H} : (\xi_k)_{k \in \mathbb{K}} \mapsto \sum_{k \in \mathbb{K}} \xi_k e_k. \quad (1.3)$$

When $\mu = \nu$ in (1.1), $(e_k)_{k \in \mathbb{K}}$ is said to be a tight frame. A simple example of a tight frame is the union of m orthonormal bases, in which case $\mu = \nu = m$. For instance, in $\mathcal{H} = L^2(\mathbb{R}^2)$, a real dual-tree wavelet decomposition is the union of two orthonormal wavelet bases [7, 30]. Curvelets [6] constitute another example of a tight frame of $L^2(\mathbb{R}^2)$. Historically, Gabor frames [16, 33] have played an important role in many inverse problems. Another common example of a frame is a Riesz basis, which corresponds to the case when $(e_k)_{k \in \mathbb{K}}$ is linearly independent or, equivalently, when F is bijective. In such instances, there exists a unique Riesz basis $(\check{e}_k)_{k \in \mathbb{K}}$ such that $(e_k)_{k \in \mathbb{K}}$ and $(\check{e}_k)_{k \in \mathbb{K}}$ are biorthogonal. Furthermore, for every $x \in \mathcal{H}$ and $(\xi_k)_{k \in \mathbb{K}} \in \ell^2(\mathbb{K})$,

$$x = F^*(\xi_k)_{k \in \mathbb{K}} \Leftrightarrow (\forall k \in \mathbb{K}) \quad \xi_k = (x | \check{e}_k). \quad (1.4)$$

When $F^{-1} = F^*$, $(e_k)_{k \in \mathbb{K}}$ is an orthonormal basis and $(\check{e}_k)_{k \in \mathbb{K}} = (e_k)_{k \in \mathbb{K}}$. Examples of Riesz bases of $L^2(\mathbb{R}^2)$ include biorthogonal bases of compactly supported dyadic wavelets having certain symmetry properties [9]. Further constructions as well as a detailed account of frame theory in Hilbert spaces can be found in [23].

The goal of the present paper is to propose a flexible convex variational framework for solving inverse problems in which *a priori* information (e.g., sparsity, distribution, statistical properties) is available about the representation of the target solution in a frame. Our analysis and our numerical algorithm will rely heavily on proximity operators. Section 2 is devoted to these operators. Our main variational formulation is presented and analysed in section 3. It consists (see problem 3.1) of minimizing the sum of a separable, possibly nondifferentiable function penalizing each coefficient of the frame decomposition individually, and a smooth function which combines other information on the problem and the data formation model. Connections with sparsity-constrained and Bayesian formulations are also established. In connection with the latter, we derive in section 4 closed-form expressions for the proximity operators associated with a variety of univariate log-concave distributions. A proximal algorithm for solving problem 3.1 is presented in section 5 and its convergence is established in infinite-dimensional spaces under mild assumptions on the penalization functions. An attractive feature of this algorithm is that it is fully split in that, at each iteration, all the functions appearing in the problem are activated individually. Finally, applications to image recovery are demonstrated in section 6.

2. Basic tool: proximity operator

2.1. Notation

Throughout, \mathcal{X} is a separable real Hilbert space with scalar product $\langle \cdot | \cdot \rangle$, norm $\|\cdot\|$ and distance d . $\Gamma_0(\mathcal{X})$ is the class of lower semicontinuous convex functions from \mathcal{X} to $]-\infty, +\infty]$ which are not identically equal to $+\infty$. The indicator function of a subset S of \mathcal{X} is

$$\iota_S : \mathcal{X} \mapsto \begin{cases} 0, & \text{if } x \in S; \\ +\infty, & \text{if } x \notin S, \end{cases} \quad (2.1)$$

its support function is $\sigma_S : \mathcal{X} \rightarrow [-\infty, +\infty] : u \mapsto \sup_{x \in S} \langle x | u \rangle$ and its distance function is $d_S : \mathcal{X} \rightarrow [0, +\infty] : x \mapsto \inf \|S - x\|$. If S is nonempty, closed and convex then,

for every $x \in \mathcal{X}$, there exists a unique point $P_S x$ in S , called the projection of x onto S , such that $\|x - P_S x\| = d_S(x)$ (further background on convex analysis will be found in [36]).

2.2. Background

Let $\varphi \in \Gamma_0(\mathcal{X})$. The subdifferential of φ at $x \in \mathcal{X}$ is the set

$$\partial\varphi(x) = \{u \in \mathcal{X} \mid (\forall y \in \mathcal{X}) \langle y - x \mid u \rangle + \varphi(x) \leq \varphi(y)\}. \tag{2.2}$$

If φ is Gâteaux differentiable at x with gradient $\nabla\varphi(x)$, then $\partial\varphi(x) = \{\nabla\varphi(x)\}$. The conjugate of φ is the function $\varphi^* \in \Gamma_0(\mathcal{X})$ defined by

$$(\forall u \in \mathcal{X}) \quad \varphi^*(u) = \sup_{x \in \mathcal{X}} \langle x \mid u \rangle - \varphi(x). \tag{2.3}$$

The continuous convex function

$$\gamma\varphi : \mathcal{X} \rightarrow \mathbb{R} : x \mapsto \inf_{y \in \mathcal{X}} \varphi(y) + \frac{1}{2\gamma} \|x - y\|^2 \tag{2.4}$$

is the Moreau envelope of index $\gamma \in]0, +\infty[$ of φ .

Definition 2.1. [28] Let $\varphi \in \Gamma_0(\mathcal{X})$. Then, for every $x \in \mathcal{X}$, the function $y \mapsto \varphi(y) + \|x - y\|^2/2$ achieves its infimum at a unique point denoted by $\text{prox}_\varphi x$. The operator $\text{prox}_\varphi : \mathcal{X} \rightarrow \mathcal{X}$ thus defined is the proximity operator of φ . Moreover,

$$(\forall x \in \mathcal{X})(\forall p \in \mathcal{X}) \quad p = \text{prox}_\varphi x \Leftrightarrow x - p \in \partial\varphi(p) \tag{2.5}$$

$$\Leftrightarrow (\forall y \in \mathcal{X}) \quad \langle y - p \mid x - p \rangle + \varphi(p) \leq \varphi(y). \tag{2.6}$$

Example 2.2. Let $\gamma \in]0, +\infty[$, let S be a nonempty convex subset of \mathcal{X} and set $\varphi = \iota_S$. Then it follows at once from (2.1), (2.4) and definition 2.1 that $\gamma\varphi = d_S^2/(2\gamma)$ and $\text{prox}_{\gamma\varphi} = P_S$.

Here are basic properties of the proximity operator.

Lemma 2.3. [15, section 2] Let $\varphi \in \Gamma_0(\mathcal{X})$. Then the following hold.

- (i) $(\forall x \in \mathcal{X}) \quad x \in \text{Argmin } \varphi \Leftrightarrow 0 \in \partial\varphi(x) \Leftrightarrow \text{prox}_\varphi x = x$.
- (ii) $(\forall x \in \mathcal{X}) \quad (\forall y \in \mathcal{X}) \|\text{prox}_\varphi x - \text{prox}_\varphi y\| \leq \|x - y\|$.
- (iii) $(\forall x \in \mathcal{X}) \quad (\forall \gamma \in]0, +\infty[) x = \text{prox}_{\gamma\varphi} x + \gamma \text{prox}_{\varphi^*/\gamma}(x/\gamma)$.

In lemma 2.3, (i) states that the minimizers of φ are characterized as the zeros of the subdifferential of φ (Fermat’s rule) or, equivalently, as the fixed points of prox_φ ; (ii) states that prox_φ is nonexpansive, which turns out to be an essential property in the convergence of iterative methods [13]; finally, (iii) is Moreau’s decomposition principle [27], which provides a powerful nonlinear decomposition rule parametrized by φ and extends in particular the standard orthogonal decomposition rule [15, remark 2.11].

Lemma 2.4. Let $\varphi \in \Gamma_0(\mathcal{X})$, let $\gamma \in]0, +\infty[$, and set $\psi = \gamma\varphi$. Then the following hold.

- (i) ψ is Fréchet-differentiable on \mathcal{X} .
- (ii) $\nabla\psi = (\text{Id} - \text{prox}_{\gamma\varphi})/\gamma = \text{prox}_{\varphi^*/\gamma}(\cdot/\gamma)$.
- (iii) $\nabla\psi$ is $(1/\gamma)$ -Lipschitz continuous.

Proof. (i) and (ii): a routine extension of [28, proposition 7.d], where $\gamma = 1$. (iii): since $\varphi^*/\gamma \in \Gamma_0(\mathcal{X})$ and (ii) asserts that $\nabla\psi = \text{prox}_{\varphi^*/\gamma}(\cdot/\gamma)$, this is a direct consequence of lemma 2.3(ii). \square

Next, we record some proximal calculus rules that will allow us to derive new proximity operators from existing ones.

Lemma 2.5. [15, lemma 2.6] *Let $\varphi \in \Gamma_0(\mathcal{X})$ and let $\mathbf{x} \in \mathcal{X}$. Then the following hold.*

- (i) *Let $\psi = \varphi + \alpha\|\cdot\|^2/2 + \langle \cdot | \mathbf{u} \rangle + \beta$, where $\mathbf{u} \in \mathcal{X}, \alpha \in [0, +\infty[$ and $\beta \in \mathbb{R}$. Then $\text{prox}_\psi \mathbf{x} = \text{prox}_{\varphi/(\alpha+1)}((\mathbf{x} - \mathbf{u})/(\alpha + 1))$.*
- (ii) *Let $\psi = \varphi(\cdot - \mathbf{z})$, where $\mathbf{z} \in \mathcal{X}$. Then $\text{prox}_\psi \mathbf{x} = \mathbf{z} + \text{prox}_\varphi(\mathbf{x} - \mathbf{z})$.*
- (iii) *Let $\psi = \varphi(\cdot/\rho)$, where $\rho \in \mathbb{R} \setminus \{0\}$. Then $\text{prox}_\psi \mathbf{x} = \rho \text{prox}_{\varphi/\rho^2}(\mathbf{x}/\rho)$.*
- (iv) *Let $\psi : \mathbf{y} \mapsto \varphi(-\mathbf{y})$. Then $\text{prox}_\psi \mathbf{x} = -\text{prox}_\varphi(-\mathbf{x})$.*

We conclude this section with some properties of proximity operators on the real line.

Lemma 2.6. *Let $\phi \in \Gamma_0(\mathbb{R})$. Then the following hold.*

- (i) [14, proposition 2.4] $\text{prox}_\phi : \mathbb{R} \rightarrow \mathbb{R}$ is increasing.
- (ii) [14, corollary 2.5] *Suppose that ϕ admits 0 as a minimizer. Then*

$$(\forall \xi \in \mathbb{R}) \begin{cases} 0 \leq \text{prox}_\phi \xi \leq \xi, & \text{if } \xi > 0; \\ \text{prox}_\phi \xi = 0, & \text{if } \xi = 0; \\ \xi \leq \text{prox}_\phi \xi \leq 0, & \text{if } \xi < 0. \end{cases} \tag{2.7}$$

This is true in particular when ϕ is even.

- (iii) [14, proposition 3.6] *Suppose that $\phi = \psi + \sigma_\Omega$, where $\psi \in \Gamma_0(\mathbb{R})$ is differentiable at 0 with $\psi'(0) = 0$, and where $\Omega \subset \mathbb{R}$ is a nonempty closed interval. Then $\text{prox}_\phi = \text{prox}_\psi \circ \text{soft}_\Omega$, where*

$$\text{soft}_\Omega = \text{prox}_{\sigma_\Omega} : \mathbb{R} \rightarrow \mathbb{R} : \xi \mapsto \begin{cases} \xi - \underline{\omega}, & \text{if } \xi < \underline{\omega}; \\ 0, & \text{if } \xi \in \Omega; \\ \xi - \bar{\omega}, & \text{if } \xi > \bar{\omega}, \end{cases} \quad \text{with} \quad \begin{cases} \underline{\omega} = \inf \Omega, \\ \bar{\omega} = \sup \Omega, \end{cases} \tag{2.8}$$

is the soft thresholder on Ω . In particular, if $\Omega = [-\omega, \omega]$ for some $\omega \in]0, +\infty[$, we obtain

$$\text{soft}_{[-\omega, \omega]} = \text{prox}_{\omega|\cdot|} : \mathbb{R} \rightarrow \mathbb{R} : \xi \mapsto \text{sign}(\xi) \max\{|\xi| - \omega, 0\}. \tag{2.9}$$

The soft-thresholding operation described in lemma 2.6(iii) is illustrated in figure 1.

2.3. Forward–backward splitting

In this section, we consider the following abstract variational framework, which will cover our main problem (problem 3.1).

Problem 2.7. Let f_1 and f_2 be functions in $\Gamma_0(\mathcal{X})$ such that f_2 is differentiable on \mathcal{X} with a β -Lipschitz continuous gradient for some $\beta \in]0, +\infty[$. The objective is to

$$\underset{\mathbf{x} \in \mathcal{X}}{\text{minimize}} \quad f_1(\mathbf{x}) + f_2(\mathbf{x}). \tag{2.10}$$




Figure 1. $\Omega = [-1, 1]$. Graphs of soft_Ω (dashed line) and prox_ϕ (solid line) in lemma 2.6(iii) with $\psi = 0.1|\cdot|^3$.

A key consequence of Fermat’s rule (lemma 2.3(i)) and (2.5) is the following characterization of the solutions to problem 2.7 which, in itself, attests the central role played by proximity operators.

Proposition 2.8. [15, proposition 3.1(iii)] *Let $\mathbf{x} \in \mathcal{X}$ and let $\gamma \in]0, +\infty[$. Then \mathbf{x} is a solution to problem 2.7 if and only if $\mathbf{x} = \text{prox}_{\gamma f_1}(\mathbf{x} - \gamma \nabla f_2(\mathbf{x}))$.*

Let $\gamma \in]0, +\infty[$ and set $T = \text{prox}_{\gamma f_1} \circ (\text{Id} - \gamma \nabla f_2)$. Proposition 2.8 asserts that a point $\mathbf{x} \in \mathcal{X}$ solves problem 2.7 if and only if $\mathbf{x} = T\mathbf{x}$. This fixed-point characterization suggests solving problem 2.7 via the successive approximation method $\mathbf{x}_{n+1} = T\mathbf{x}_n$, for suitable values of the ‘step size’ parameter γ . The next result describes an algorithm in this vein, which is based on the forward–backward splitting method for monotone operators [13]. It allows for inexact evaluations of the operators prox_{f_1} and ∇f_2 via the incorporation of the error sequences $(\mathbf{a}_n)_{n \in \mathbb{N}}$ and $(\mathbf{b}_n)_{n \in \mathbb{N}}$, respectively, as well as for iteration-dependent relaxation parameters $(\lambda_n)_{n \in \mathbb{N}}$ and step sizes $(\gamma_n)_{n \in \mathbb{N}}$.

Theorem 2.9. [15, theorem 3.4(i)] *Suppose that $\text{Argmin}(f_1 + f_2) \neq \emptyset$. Let $(\gamma_n)_{n \in \mathbb{N}}$ be a sequence in $]0, +\infty[$ such that $0 < \inf_{n \in \mathbb{N}} \gamma_n \leq \sup_{n \in \mathbb{N}} \gamma_n < 2/\beta$, let $(\lambda_n)_{n \in \mathbb{N}}$ be a sequence in $]0, 1[$ such that $\inf_{n \in \mathbb{N}} \lambda_n > 0$, and let $(\mathbf{a}_n)_{n \in \mathbb{N}}$ and $(\mathbf{b}_n)_{n \in \mathbb{N}}$ be sequences in \mathcal{X} such that $\sum_{n \in \mathbb{N}} \|\mathbf{a}_n\| < +\infty$ and $\sum_{n \in \mathbb{N}} \|\mathbf{b}_n\| < +\infty$. Fix $\mathbf{x}_0 \in \mathcal{X}$ and, for every $n \in \mathbb{N}$, set*

$$\mathbf{x}_{n+1} = \mathbf{x}_n + \lambda_n (\text{prox}_{\gamma_n f_1}(\mathbf{x}_n - \gamma_n (\nabla f_2(\mathbf{x}_n) + \mathbf{b}_n)) + \mathbf{a}_n - \mathbf{x}_n). \tag{2.11}$$

Then $(\mathbf{x}_n)_{n \in \mathbb{N}}$ converges weakly to a solution to problem 2.7.

2.4. Decomposition formula

The following decomposition property, which extends [15, example 2.19], will be instrumental in our analysis.

Proposition 2.10. Set $\Upsilon : \mathcal{X} \rightarrow]-\infty, +\infty] : x \mapsto \sum_{i \in \mathbb{I}} \psi_i(\langle x | o_i \rangle)$, where:

- (i) $\emptyset \neq \mathbb{I} \subset \mathbb{N}$;
- (ii) $(o_i)_{i \in \mathbb{I}}$ is an orthonormal basis of \mathcal{X} ;
- (iii) $(\psi_i)_{i \in \mathbb{I}}$ are functions in $\Gamma_0(\mathbb{R})$;
- (iv) Either \mathbb{I} is finite, or there exists a subset \mathbb{J} of \mathbb{I} such that:
 - (a) $\mathbb{I} \setminus \mathbb{J}$ is finite;
 - (b) $(\forall i \in \mathbb{J}) \psi_i \geq 0$;
 - (c) there exists a sequence $(\zeta_i)_{i \in \mathbb{J}}$ in \mathbb{R} such that $\sum_{i \in \mathbb{J}} |\zeta_i|^2 < +\infty$, $\sum_{i \in \mathbb{J}} |\text{prox}_{\psi_i} \zeta_i|^2 < +\infty$, and $\sum_{i \in \mathbb{J}} \psi_i(\zeta_i) < +\infty$.

Then $\Upsilon \in \Gamma_0(\mathcal{X})$ and $(\forall x \in \mathcal{X}) \text{prox}_{\Upsilon} x = \sum_{i \in \mathbb{I}} (\text{prox}_{\psi_i} \langle x | o_i \rangle) o_i$.

Proof. We treat only the case when \mathbb{I} is infinite as the case when \mathbb{I} is finite will follow trivially the arguments presented below. Fix, for every $i \in \mathbb{I} \setminus \mathbb{J}$, $\zeta_i \in \mathbb{R}$ such that $\psi_i(\zeta_i) < +\infty$ and set $z = \sum_{i \in \mathbb{I}} \zeta_i o_i$. Then (iv) implies that $\sum_{i \in \mathbb{I}} \zeta_i^2 < +\infty$ and, in view of (ii), that $z \in \mathcal{X}$. Moreover, $\Upsilon(z) = \sum_{i \in \mathbb{J}} \psi_i(\zeta_i) + \sum_{i \in \mathbb{I} \setminus \mathbb{J}} \psi_i(\zeta_i) < +\infty$.

Let us show that $\Upsilon \in \Gamma_0(\mathcal{X})$. As just seen, $\Upsilon(z) < +\infty$ and, therefore, $\Upsilon \neq +\infty$. Next, we observe that, by virtue of (iii), the functions $(\psi_i(\langle \cdot | o_i \rangle))_{i \in \mathbb{I}}$ are lower semicontinuous and convex. As a result, $\sum_{i \in \mathbb{I} \setminus \mathbb{J}} \psi_i(\langle \cdot | o_i \rangle)$ is lower semicontinuous and convex, as a finite sum of such functions. Thus, to show that $\Upsilon \in \Gamma_0(\mathcal{X})$, it remains to show that $\Upsilon_{\mathbb{J}} = \sum_{i \in \mathbb{J}} \psi_i(\langle \cdot | o_i \rangle)$ is lower semicontinuous and convex. It follows from (iv)(b) that

$$\Upsilon_{\mathbb{J}} = \sup_{\substack{\mathbb{J}' \subset \mathbb{J} \\ \mathbb{J}' \text{ finite}}} \sum_{i \in \mathbb{J}'} \psi_i(\langle \cdot | o_i \rangle). \tag{2.12}$$

However, as above, each finite sum $\sum_{i \in \mathbb{J}'} \psi_i(\langle \cdot | o_i \rangle)$ is lower semicontinuous and convex. Therefore, $\Upsilon_{\mathbb{J}}$ is likewise as the supremum of a family of lower semicontinuous convex functions.

Now fix $x \in \mathcal{X}$ and set

$$(\forall i \in \mathbb{I}) \quad \xi_i = \langle x | o_i \rangle \quad \text{and} \quad \pi_i = \text{prox}_{\psi_i} \xi_i. \tag{2.13}$$

It follows from (iv)(a) and (iv)(c) that

$$\sum_{i \in \mathbb{I}} |\text{prox}_{\psi_i} \zeta_i|^2 = \sum_{i \in \mathbb{J}} |\text{prox}_{\psi_i} \zeta_i|^2 + \sum_{i \in \mathbb{I} \setminus \mathbb{J}} |\text{prox}_{\psi_i} \zeta_i|^2 < +\infty. \tag{2.14}$$

Hence, we derive from lemma 2.3(ii) and (ii) that

$$\begin{aligned} \frac{1}{2} \sum_{i \in \mathbb{I}} |\pi_i|^2 &\leq \sum_{i \in \mathbb{I}} |\pi_i - \text{prox}_{\psi_i} \zeta_i|^2 + \sum_{i \in \mathbb{I}} |\text{prox}_{\psi_i} \zeta_i|^2 \\ &= \sum_{i \in \mathbb{I}} |\text{prox}_{\psi_i} \xi_i - \text{prox}_{\psi_i} \zeta_i|^2 + \sum_{i \in \mathbb{I}} |\text{prox}_{\psi_i} \zeta_i|^2 \\ &\leq \sum_{i \in \mathbb{I}} |\xi_i - \zeta_i|^2 + \sum_{i \in \mathbb{I}} |\text{prox}_{\psi_i} \zeta_i|^2 \\ &= \|x - z\|^2 + \sum_{i \in \mathbb{I}} |\text{prox}_{\psi_i} \zeta_i|^2 \\ &< +\infty. \end{aligned} \tag{2.15}$$

Let us set $p = \sum_{i \in \mathbb{I}} \pi_i o_i$. Then it follows from (2.15) and (ii) that $p \in \mathcal{X}$. On the other hand, we derive from (2.13) and (2.6) that

$$(\forall i \in \mathbb{I})(\forall \eta \in \mathbb{R}) \quad (\eta - \pi_i)(\xi_i - \pi_i) + \psi_i(\pi_i) \leq \psi_i(\eta). \tag{2.16}$$

Hence, by Parseval and (ii),

$$\begin{aligned}
 (\forall y \in \mathcal{X}) \quad \langle y - p \mid x - p \rangle + \Upsilon(p) &= \sum_{i \in \mathbb{I}} \langle y - p \mid o_i \rangle \langle x - p \mid o_i \rangle + \sum_{i \in \mathbb{I}} \psi_i(\pi_i) \\
 &= \sum_{i \in \mathbb{I}} (\langle y \mid o_i \rangle - \pi_i)(\xi_i - \pi_i) + \psi_i(\pi_i) \\
 &\leq \sum_{i \in \mathbb{I}} \psi_i(\langle y \mid o_i \rangle) \\
 &= \Upsilon(y).
 \end{aligned} \tag{2.17}$$

Invoking (2.6) once again, we conclude that $p = \text{prox}_{\Upsilon} x$. □

3. Problem formulation

3.1. Assumptions and problem statement

Throughout, \mathcal{H} is a separable real Hilbert space with scalar product $\langle \cdot \mid \cdot \rangle$, norm $\| \cdot \|$ and distance d . The index set \mathbb{K} is either $\{1, \dots, K\}$ ($K \in \mathbb{N}$) or \mathbb{N} , according as \mathcal{H} is finite or infinite dimensional. Moreover, $(e_k)_{k \in \mathbb{K}}$ is a frame in \mathcal{H} with constants μ and ν (see (1.1)) and frame operator F (see (1.2)). Finally, the sequence of frame coefficients of a generic point $x \in \mathcal{H}$ will be denoted by \mathbf{x} , i.e., $\mathbf{x} = (\xi_k)_{k \in \mathbb{K}}$, where $x = \sum_{k \in \mathbb{K}} \xi_k e_k$.

Let $\bar{x} \in \mathcal{H}$ be the target solution of the underlying inverse problem. Our basic premise is that *a priori* information is available about the coefficients $(\bar{\xi}_k)_{k \in \mathbb{K}}$ of the decomposition

$$\bar{x} = \sum_{k \in \mathbb{K}} \bar{\xi}_k e_k \tag{3.1}$$

of \bar{x} in $(e_k)_{k \in \mathbb{K}}$. To recover \bar{x} , it is therefore natural to formulate a variational problem in the space $\ell^2(\mathbb{K})$ of frame coefficients, where *a priori* information on $(\bar{\xi}_k)_{k \in \mathbb{K}}$ can be easily incorporated. More precisely, a solution will assume the form $\tilde{x} = \sum_{k \in \mathbb{K}} \tilde{\xi}_k e_k$, where $(\tilde{\xi}_k)_{k \in \mathbb{K}}$ is a solution to the following problem.

Problem 3.1. Let $(\phi_k)_{k \in \mathbb{K}}$ be functions in $\Gamma_0(\mathbb{R})$ such that either $\mathbb{K} = \{1, \dots, K\}$ with $K \in \mathbb{N}$, or $\mathbb{K} = \mathbb{N}$ and there exists a subset \mathbb{L} of \mathbb{K} such that

- (i) $\mathbb{K} \setminus \mathbb{L}$ is finite;
- (ii) $(\forall k \in \mathbb{L}) \phi_k \geq 0$;
- (iii) there exists a sequence $(\zeta_k)_{k \in \mathbb{L}}$ in \mathbb{R} such that $\sum_{k \in \mathbb{L}} |\zeta_k|^2 < +\infty$, $\sum_{k \in \mathbb{L}} |\text{prox}_{\phi_k} \zeta_k|^2 < +\infty$ and $\sum_{k \in \mathbb{L}} \phi_k(\zeta_k) < +\infty$.

In addition, let $\Psi \in \Gamma_0(\mathcal{H})$ be differentiable on \mathcal{H} with a τ -Lipschitz continuous gradient for some $\tau \in]0, +\infty[$. The objective is to

$$\underset{(\xi_k)_{k \in \mathbb{K}} \in \ell^2(\mathbb{K})}{\text{minimize}} \sum_{k \in \mathbb{K}} \phi_k(\xi_k) + \Psi \left(\sum_{k \in \mathbb{K}} \xi_k e_k \right). \tag{3.2}$$

Remark 3.2.

- (i) The functions $(\phi_k)_{k \in \mathbb{K}}$ in problem 3.1 need not be differentiable. As will be seen in section 3.3.1, this feature is essential in sparsity-constrained problems.
- (ii) Suppose that $\mathbb{K} = \mathbb{N}$. Then conditions (ii) and (iii) in problem 3.1 hold when, for every $k \in \mathbb{L}$, ϕ_k admits a minimizer ζ_k such that $\phi_k(\zeta_k) = 0$ and $\sum_{k \in \mathbb{L}} |\zeta_k|^2 < +\infty$. Indeed, lemma 2.3(i) yields $\sum_{k \in \mathbb{L}} |\text{prox}_{\phi_k} \zeta_k|^2 = \sum_{k \in \mathbb{L}} |\zeta_k|^2 < +\infty$ and $\sum_{k \in \mathbb{L}} \phi_k(\zeta_k) = 0$. In particular, conditions (ii) and (iii) in problem 3.1 hold when $(\forall k \in \mathbb{L}) \phi_k \geq \phi_k(0) = 0$, which amounts to setting $\zeta_k \equiv 0$.

3.2. Existence and characterization of solutions

We first address the issue of the existence of solutions to problem 3.1. Recall that a function $\varphi : \mathcal{H} \rightarrow]-\infty, +\infty]$ is said to be coercive if $\lim_{\|x\| \rightarrow +\infty} \varphi(x) = +\infty$.

Proposition 3.3. *Suppose that one of the following holds.*

- (i) *The function $(\xi_k)_{k \in \mathbb{K}} \mapsto \sum_{k \in \mathbb{K}} \phi_k(\xi_k) + \Psi(F^*(\xi_k)_{k \in \mathbb{K}})$ is coercive.*
- (ii) *$\inf_{k \in \mathbb{K}} \inf \phi_k(\mathbb{R}) > -\infty$, Ψ is coercive, and $(e_k)_{k \in \mathbb{K}}$ is a Riesz basis.*
- (iii) *$\inf \Psi(\mathcal{H}) > -\infty$ and one of the following properties is satisfied.*
 - (a) *The function $(\xi_k)_{k \in \mathbb{K}} \mapsto \sum_{k \in \mathbb{K}} \phi_k(\xi_k)$ is coercive.*
 - (b) *There exist $\omega \in]0, +\infty[$ and $p \in [1, 2]$ such that $(\forall k \in \mathbb{K}) \phi_k \geq \omega \cdot |\cdot|^p$.*
 - (c) *\mathbb{K} is finite and the functions $(\phi_k)_{k \in \mathbb{K}}$ are coercive.*

Then problem 3.1 admits a solution.

Proof. We denote by $\mathbf{x} = (\xi_k)_{k \in \mathbb{K}}$ a generic element in $\ell^2(\mathbb{K})$ and by $\|\mathbf{x}\| = \sqrt{\sum_{k \in \mathbb{K}} |\xi_k|^2}$ its norm. Set

$$f_1 : \mathbf{x} \mapsto \sum_{k \in \mathbb{K}} \phi_k(\xi_k) \quad \text{and} \quad f_2 = \Psi \circ F^*. \tag{3.3}$$

First, suppose that (i) holds. Then it follows from the assumptions on $(\phi_k)_{k \in \mathbb{K}}$ in problem 3.1 and proposition 2.10 that $f_1 \in \Gamma_0(\ell^2(\mathbb{K}))$. On the other hand, since Ψ is a finite function in $\Gamma_0(\mathcal{H})$ and $F^* : \ell^2(\mathbb{K}) \rightarrow \mathcal{H}$ is linear and bounded, f_2 is a finite function in $\Gamma_0(\ell^2(\mathbb{K}))$. Altogether, $f_1 + f_2 \in \Gamma_0(\ell^2(\mathbb{K}))$ and the claim follows from [36, theorem 2.5.1(ii)].

Next, suppose that (ii) holds. In view of (i), since f_1 is bounded below, it is enough to show that f_2 is coercive. Since $(e_k)_{k \in \mathbb{K}}$ is a Riesz basis, we have [26]

$$(\forall \mathbf{x} \in \ell^2(\mathbb{K})) \quad \|F^*\mathbf{x}\| \geq \sqrt{\mu}\|\mathbf{x}\|. \tag{3.4}$$

In turn, the coercivity of Ψ implies that $\lim_{\|\mathbf{x}\| \rightarrow +\infty} \|\Psi(F^*\mathbf{x})\| = +\infty$.

Now, suppose that (iii) holds. In case (iii)(a), since Ψ is bounded below, f_2 is likewise. In turn, the coercivity of f_1 implies that of $f_1 + f_2$, hence the result by (i). Now suppose that (iii)(b) is satisfied and let $\mathbf{x} \in \ell^2(\mathbb{K})$. Then

$$f_1(\mathbf{x}) = \sum_{k \in \mathbb{K}} \phi_k(\xi_k) \geq \omega \sum_{k \in \mathbb{K}} |\xi_k|^p \geq \omega \|\mathbf{x}\|^p. \tag{3.5}$$

Therefore f_1 is coercive and the claim follows from (iii)(a). Finally, suppose that (iii)(c) is satisfied. In view of (iii)(a), it is enough to show that f_1 is coercive. To this end, fix $\rho \in]0, +\infty[$ and recall that $\mathbb{K} = \{1, \dots, K\}$. Let us set $\lambda = \min_{k \in \mathbb{K}} \inf \phi_k(\mathbb{R})$. Since the functions $(\phi_k)_{k \in \mathbb{K}}$ are coercive and in $\Gamma_0(\mathbb{R})$, it follows from [36, theorem 2.5.1(ii)] that $\lambda \in \mathbb{R}$. Coercivity also implies that we can find $\delta \in]0, +\infty[$ such that

$$(\forall \xi \in \mathbb{R}) \quad |\xi| \geq \delta/\sqrt{K} \quad \Rightarrow \quad \min_{k \in \mathbb{K}} \phi_k(\xi) \geq \rho + (1 - K)\lambda. \tag{3.6}$$

Now take $\mathbf{x} \in \ell^2(\mathbb{K})$ such that $\|\mathbf{x}\| \geq \delta$ and fix $\ell \in \mathbb{K}$ such that $|\xi_\ell| = \max_{k \in \mathbb{K}} |\xi_k|$. Then $|\xi_\ell| \geq \delta/\sqrt{K}$ and therefore (3.6) yields

$$f_1(\mathbf{x}) = \sum_{k \in \mathbb{K}} \phi_k(\xi_k) \geq \rho + (1 - K)\lambda + \sum_{k \in \mathbb{K} \setminus \{\ell\}} \phi_k(\xi_k) \geq \rho + (1 - K)\lambda + (K - 1)\lambda = \rho, \tag{3.7}$$

which shows that f_1 is coercive. □

Next, we turn our attention to the characterization of the solutions to problem 3.1.

Proposition 3.4. *Let $(\xi_k)_{k \in \mathbb{K}} \in \ell^2(\mathbb{K})$, let $(\eta_k)_{k \in \mathbb{K}} = (F \circ \nabla \Psi \circ F^*)(\xi_k)_{k \in \mathbb{K}}$ and let $\gamma \in]0, +\infty[$. Then $(\xi_k)_{k \in \mathbb{K}}$ solves problem 3.1 if and only if $(\forall k \in \mathbb{K}) \xi_k = \text{prox}_{\gamma \phi_k}(\xi_k - \gamma \eta_k)$.*

Proof. Set $\mathcal{X} = \ell^2(\mathbb{K})$ and let $(o_k)_{k \in \mathbb{K}}$ denote the canonical orthonormal basis of $\ell^2(\mathbb{K})$. Then (3.2) can be written as

$$\underset{x \in \mathcal{X}}{\text{minimize}} \sum_{k \in \mathbb{K}} \phi_k(\langle x \mid o_k \rangle) + \Psi(F^*x). \tag{3.8}$$

Now set

$$f_1 = \sum_{k \in \mathbb{K}} \phi_k(\langle \cdot \mid o_k \rangle) \quad \text{and} \quad f_2 = \Psi \circ F^*. \tag{3.9}$$

Then, in the light of the assumptions on $(\phi_k)_{k \in \mathbb{K}}$ in problem 3.1, proposition 2.10 yields $f_1 \in \Gamma_0(\mathcal{X})$. On the other hand, since Ψ is a finite function in $\Gamma_0(\mathcal{H})$ and $F^* : \mathcal{X} \rightarrow \mathcal{H}$ is linear and bounded, we have $f_2 \in \Gamma_0(\mathcal{X})$. In addition, since $\nabla \Psi$ is Lipschitz continuous, so is $\nabla f_2 = F \circ \nabla \Psi \circ F^*$. Altogether, (3.8) conforms to the format of problem 2.7. Now set $x = (\xi_k)_{k \in \mathbb{K}}$. Then it follows from proposition 2.8, (3.9) and proposition 2.10 that

$$\begin{aligned} (\xi_k)_{k \in \mathbb{K}} \text{ solves problem 3.1} &\Leftrightarrow x = \text{prox}_{\gamma f_1}(x - \gamma \nabla f_2(x)) \\ &\Leftrightarrow x = \text{prox}_{\gamma f_1}(x - \gamma (F \circ \nabla \Psi \circ F^*)(x)) \\ &\Leftrightarrow (\xi_k)_{k \in \mathbb{K}} = \text{prox}_{\gamma f_1}(\xi_k - \gamma \eta_k)_{k \in \mathbb{K}} \\ &\Leftrightarrow (\xi_k)_{k \in \mathbb{K}} = (\text{prox}_{\gamma \phi_k}(\xi_k - \gamma \eta_k))_{k \in \mathbb{K}}, \end{aligned} \tag{3.10}$$

which provides the desired characterization. □

3.3. Specific frameworks

In problem 3.1, the functions $(\phi_k)_{k \in \mathbb{K}}$ penalize the frame coefficients $(\xi_k)_{k \in \mathbb{K}}$, while the function Ψ penalizes $x = F^*(\xi_k)_{k \in \mathbb{K}} = \sum_{k \in \mathbb{K}} \xi_k e_k$, thereby modelling direct constraints on \bar{x} . This flexible framework makes it possible to model a wide range of inverse problems. Two important instances are presented below.

3.3.1. Inverse problems with sparsity constraints. A common objective in selecting the frame $(e_k)_{k \in \mathbb{K}}$ is to obtain a sparse representation of the target solution \bar{x} in the sense that most of the coefficients $(\bar{\xi}_k)_{k \in \mathbb{K}}$ in (3.1) are zero. By choosing $\phi_k = \omega_k |\cdot|$ with $\omega_k > 0$ in problem 3.1, one aims at setting to zero the k th coefficient if it falls into the interval $[-\omega_k, \omega_k]$, hence promoting sparsity (see [17, 22, 34] for special cases). Note that in this case, it follows from proposition 3.4 and (2.9) that a solution $(\xi_k)_{k \in \mathbb{K}}$ to problem 3.1 is characterized by the soft thresholding identities (see also figure 1)

$$(\forall k \in \mathbb{K}) \quad \xi_k = \text{prox}_{\omega_k |\cdot|}(\xi_k - \eta_k) = \text{soft}_{[-\omega_k, \omega_k]}(\xi_k - \eta_k), \tag{3.11}$$

where $(\eta_k)_{k \in \mathbb{K}} = (F \circ \nabla \Psi \circ F^*)(\xi_k)_{k \in \mathbb{K}}$. More generally, to aim at zeroing a coefficient falling into a closed interval $\Omega_k \subset \mathbb{R}$, one can use the function $\phi_k = \psi_k + \sigma_{\Omega_k}$, where ψ_k satisfies $0 = \psi_k(0) \leq \psi_k \in \Gamma_0(\mathbb{R})$ and is differentiable at 0 [14, proposition 3.2]. This construct actually characterizes all thresholders on Ω_k that have properties suitable for their use in iterative methods [14, theorem 3.3]. A decomposition rule for computing the resulting thresholders is supplied in lemma 2.6(iii).

Let us now discuss possible choices for the smooth function Ψ . Suppose that the problem under consideration is to recover $\bar{x} \in \mathcal{H}$ from q observations

$$z_i = T_i \bar{x} + v_i, \quad 1 \leq i \leq q, \tag{3.12}$$

where T_i is a bounded linear operator from \mathcal{H} to a real Hilbert space \mathcal{G}_i , $z_i \in \mathcal{G}_i$, and $v_i \in \mathcal{G}_i$ is the realization of a noise process. A standard data fidelity criterion in such instances is the function $x \mapsto \sum_{i=1}^q \alpha_i \|T_i x - z_i\|^2$, where $(\alpha_i)_{1 \leq i \leq q}$ are strictly positive reals, see e.g., [12, 21]. In addition, assume that *a priori* information is available that constrains \bar{x} to lie in some closed convex subsets $(S_i)_{1 \leq i \leq m}$ of \mathcal{H} (see [10, 31] and the references therein for example). These constraints can be aggregated via the cost function $x \mapsto \sum_{i=1}^m \vartheta_i d_{S_i}^2(x)$, where $(\vartheta_i)_{1 \leq i \leq m}$ are strictly positive reals [5, 11]. These two objectives can be combined by using the function

$$\Psi : x \mapsto \frac{1}{2} \sum_{i=1}^q \alpha_i \|T_i x - z_i\|^2 + \frac{1}{2} \sum_{i=1}^m \vartheta_i d_{S_i}^2(x) \tag{3.13}$$

in problem 3.1. This function is indeed differentiable and its gradient

$$\nabla \Psi : x \mapsto \sum_{i=1}^q \alpha_i T_i^* (T_i x - z_i) + \sum_{i=1}^m \vartheta_i (x - P_{S_i} x) \tag{3.14}$$

has Lipschitz constant [14, section 5.1]

$$\tau = \left\| \sum_{i=1}^q \alpha_i T_i^* T_i \right\| + \sum_{i=1}^m \vartheta_i. \tag{3.15}$$

In instances when $\left\| \sum_{i=1}^q \alpha_i T_i^* T_i \right\|$ cannot be evaluated directly, it can be majorized by $\sum_{i=1}^q \alpha_i \|T_i\|^2$. It should be noted that, more generally, lemma 2.4(iii) implies that Ψ remains Lipschitz continuous if the term $\sum_{i=1}^m \vartheta_i d_{S_i}^2(x)$ in (3.13) is replaced by a sum of Moreau envelopes (see [13, section 6.3] and [15, section 4.1] for related frameworks).

3.3.2. Bayesian statistical framework. A standard linear inverse problem is to recover $\bar{x} \in \mathcal{H}$ from an observation

$$z = T\bar{x} + v, \tag{3.16}$$

in a real Hilbert space \mathcal{G} , where $T : \mathcal{H} \rightarrow \mathcal{G}$ is a bounded linear operator and where $v \in \mathcal{G}$ is the realization of an additive noise perturbation. If $\bar{x} = (\bar{x}_k)_{k \in \mathbb{K}}$ denotes the coefficients of \bar{x} in $(e_k)_{k \in \mathbb{K}}$, (3.16) can be written as

$$z = TF^*\bar{x} + v. \tag{3.17}$$

For the sake of simplicity, the following assumptions regarding (3.17) are made in this section (with the usual convention $\ln 0 = -\infty$).

Assumption 3.5.

- (i) $\mathcal{H} = \mathbb{R}^N$, $\mathcal{G} = \mathbb{R}^M$ and $\mathbb{K} = \{1, \dots, K\}$, where $K \geq N$.
- (ii) The vectors \bar{x} , z and v are, respectively, realizations of real-valued random vectors \bar{X} , Z and V defined on the same probability space.
- (iii) The random vectors \bar{X} and V are mutually independent and have probability density functions $f_{\bar{x}}$ and f_v , respectively.
- (iv) The components of \bar{X} are independent with upper-semicontinuous log-concave densities.
- (v) The function $\ln f_v$ is concave and differentiable with a Lipschitz continuous gradient.

Under assumption 3.5, a common Bayesian approach for estimating \bar{x} from z consists in applying a maximum *a posteriori* (MAP) rule [3, 4, 32], which amounts to maximizing the posterior probability density $f_{\bar{x}|Z=z}$. Thus, \tilde{x} is a MAP estimate of \bar{x} if

$$(\forall x \in \mathbb{R}^K) \quad f_{\bar{x}|Z=z}(\tilde{x}) \geq f_{\bar{x}|Z=z}(x). \tag{3.18}$$

Using Bayes' formula, this amounts to solving

$$\underset{x \in \mathbb{R}^K}{\text{minimize}} - \ln f_{\bar{X}}(x) - \ln f_{z|\bar{X}=x}(z). \tag{3.19}$$

In view of (3.17), this is also equivalent to solving

$$\underset{x \in \mathbb{R}^K}{\text{minimize}} - \ln f_{\bar{X}}(x) - \ln f_V(z - TF^*x). \tag{3.20}$$

Under assumption 3.5, this convex optimization problem is a special case of problem 3.1. Indeed, assumption 3.5(iv) allows us to write, without loss of generality, the prior density as

$$(\forall (\xi_k)_{k \in \mathbb{K}} \in \mathbb{R}^K) \quad f_{\bar{X}}((\xi_k)_{k \in \mathbb{K}}) \propto \prod_{k=1}^K \exp(-\phi_k(\xi_k)), \tag{3.21}$$

where $(\phi_k)_{k \in \mathbb{K}}$ are the so-called potential functions of the marginal probability density functions of \bar{X} . It also follows from assumption 3.5(iv) that the functions $(\phi_k)_{k \in \mathbb{K}}$ are in $\Gamma_0(\mathbb{R})$. Now set

$$(\forall x \in \mathcal{H}) \quad \Psi(x) = - \ln f_V(z - Tx). \tag{3.22}$$

Then assumption 3.5(v) asserts that $\Psi \in \Gamma_0(\mathcal{H})$ is differentiable with a Lipschitz continuous gradient. Altogether, (3.20) reduces to problem 3.1.

Remark 3.6. In the simple case when V is a zero-mean Gaussian vector with an invertible covariance matrix Λ , the function Ψ reduces (up to an additive constant) to the residual energy function $x \mapsto \langle \Lambda^{-1}(z - Tx) | z - Tx \rangle / 2$. When \bar{X} is further assumed to be Gaussian, the solution to problem 3.1 is a linear function of z . Recall that the MAP estimate coincides with the minimum mean-square error estimate under such Gaussian models for both V and \bar{X} [35, section 2.4].

Remark 3.7. An alternative Bayesian strategy would be to determine a MAP estimate of \bar{x} . This would lead to

$$\underset{x \in \mathcal{H}}{\text{minimize}} - \ln f_{\bar{X}}(x) - \ln f_V(z - Tx), \tag{3.23}$$

where $f_{\bar{X}}$ can be deduced from (3.21) through the change of variable $\bar{X} = F^*X$. In the case of an orthonormal basis decomposition, it is easy to check that (3.23) is equivalent to problem (3.20). By contrast, when F corresponds to an overcomplete frame, the expression of $f_{\bar{X}}$ becomes involved and (3.23) is usually much less tractable than problem 3.1. As will be seen in section 5, the latter can be solved via a simple splitting algorithm.

Remark 3.8. Let us decompose the observation vector as $z = [z_1^\top, \dots, z_q^\top]^\top$ and the matrix representing T as $[T_1^\top, \dots, T_q^\top]^\top$ where, for every $i \in \{1, \dots, q\}$, $z_i \in \mathbb{R}^{M_i}$ and $T_i \in \mathbb{R}^{M_i \times N}$ with $\sum_{i=1}^q M_i = M$. Furthermore, assume that V is a zero-mean Gaussian vector with the diagonal covariance matrix

$$\Lambda = \begin{bmatrix} \alpha_1^{-1} I_{M_1} & 0 & \dots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & \alpha_q^{-1} I_{M_q} \end{bmatrix}, \tag{3.24}$$

where $(\alpha_i)_{1 \leq i \leq q}$ are strictly positive reals and I_{M_i} , $1 \leq i \leq q$, is the identity matrix of size $M_i \times M_i$. Then Ψ reduces to the first term in (3.13) where $\mathcal{G}_i = \mathbb{R}^{M_i}$ and the MAP estimation problem under assumption 3.5 becomes a special case of the problem addressed in section 3.3.1 with $m = 0$.

4. Proximity operators associated with log-concave densities

As discussed in section 3.3.2, the functions $(\phi_k)_{k \in \mathbb{K}}$ in (3.2) act as the potential functions of log-concave univariate probability densities modelling the frame coefficients individually in Bayesian formulations. On the other hand, the proximity operators of such functions will, via proposition 2.10, play a central role in section 5. Hereafter, we derive closed-form expressions for these proximity operators in the case of some classical log-concave univariate probability densities [19, chapters VII and IX].

Let us start with a few observations.

Remark 4.1. Let $\phi \in \Gamma_0(\mathbb{R})$.

- (i) It follows from definition 2.1 that $(\forall \xi \in \mathbb{R}) \phi(\text{prox}_\phi \xi) < +\infty$.
- (ii) If ϕ is even, then it follows from lemma 2.5(iv) that prox_ϕ is odd. Therefore, in such instances, it will be enough to determine $\text{prox}_\phi \xi$ for $\xi \geq 0$ and to extend the result to $\xi < 0$ by antisymmetry.
- (iii) Let $\xi \in \mathbb{R}$. If ϕ is differentiable at $\text{prox}_\phi \xi$, then (2.5) yields

$$(\forall \pi \in \mathbb{R}) \quad \pi = \text{prox}_\phi \xi \quad \Leftrightarrow \quad \pi + \phi'(\pi) = \xi. \quad (4.1)$$

We now examine some concrete examples.

Example 4.2 (Laplace distribution). Let $\omega \in]0, +\infty[$ and set

$$\phi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \omega|\xi|. \quad (4.2)$$

Then, for every $\xi \in \mathbb{R}$, $\text{prox}_\phi \xi = \text{soft}_{[-\omega, \omega]} \xi = \text{sign}(\xi) \max\{|\xi| - \omega, 0\}$.

Proof. Apply lemma 2.6(iii) with $\psi = 0$ and $\Omega = [-\omega, \omega]$. □

Example 4.3 (Gaussian distribution). Let $\tau \in]0, +\infty[$ and set

$$\phi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \tau|\xi|^2. \quad (4.3)$$

Then, for every $\xi \in \mathbb{R}$, $\text{prox}_\phi \xi = \xi/(2\tau + 1)$.

Proof. Apply lemma 2.5(i) with $\mathcal{X} = \mathbb{R}$, $\varphi = 0$, $\alpha = 2\tau$ and $\mathbf{u} = 0$. □

Example 4.4 (generalized Gaussian distribution). Let $p \in]1, +\infty[$, $\kappa \in]0, +\infty[$ and set

$$\phi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \kappa|\xi|^p. \quad (4.4)$$

Then, for every $\xi \in \mathbb{R}$, $\text{prox}_\phi \xi = \text{sign}(\xi)\varrho$ where ϱ is the unique solution in $[0, +\infty[$ to

$$\varrho + p\kappa\varrho^{p-1} = |\xi|. \quad (4.5)$$

In particular, the following hold:

- (i) $\text{prox}_\phi \xi = \xi + \frac{4\kappa}{3 \times 2^{1/3}} ((\chi - \xi)^{1/3} - (\chi + \xi)^{1/3})$, where $\chi = \sqrt{\xi^2 + 256\kappa^3/729}$, if $p = 4/3$;
- (ii) $\text{prox}_\phi \xi = \xi + 9\kappa^2 \text{sign}(\xi)(1 - \sqrt{1 + 16|\xi|/(9\kappa^2)})/8$, if $p = 3/2$;
- (iii) $\text{prox}_\phi \xi = \text{sign}(\xi)(\sqrt{1 + 12\kappa|\xi|} - 1)/(6\kappa)$, if $p = 3$;
- (iv) $\text{prox}_\phi \xi = \left(\frac{\chi + \xi}{8\kappa}\right)^{1/3} - \left(\frac{\chi - \xi}{8\kappa}\right)^{1/3}$, where $\chi = \sqrt{\xi^2 + 1/(27\kappa)}$, if $p = 4$.

Proof. Let $\xi \in \mathbb{R}$ and set $\pi = \text{prox}_\phi \xi$. As seen in remark 4.1(ii), because ϕ is even, it is enough to assume that $\xi \geq 0$. Since ϕ is differentiable, it follows from (2.7) and (4.1) that π is the unique solution in $[0, +\infty[$ to

$$\pi + p\kappa\pi^{p-1} = \xi, \tag{4.6}$$

which provides (4.5). For $p = 3$, π is the solution in $[0, +\infty[$ to the equation $\pi + 3\kappa\pi^2 - \xi = 0$, i.e., $\pi = (\sqrt{1 + 12\kappa\xi} - 1)/(6\kappa)$ and we obtain (iv) by antisymmetry. In turn, since $(2|\cdot|^{3/2}/3)^* = |\cdot|^3/3$, lemma 2.3(iii) with $\gamma = 3\kappa/2$ yields $\pi = \text{prox}_{\gamma(2|\cdot|^{3/2}/3)}\xi = \xi - \gamma \text{prox}_{(3\gamma)^{-1}|\cdot|^3}(\xi/\gamma) = \xi + 9\kappa^2 \text{sign}(\xi)(1 - \sqrt{1 + 16|\xi|/(9\kappa^2)})/8$, which proves (ii). Now, let $p = 4$. Then (4.6) asserts that π is the unique solution in $[0, +\infty[$ to the third degree equation $4\kappa\pi^3 + \pi - \xi = 0$, namely $\pi = (\sqrt{\alpha^2 + \beta^3} - \alpha)^{1/3} - (\sqrt{\alpha^2 + \beta^3} + \alpha)^{1/3}$, where $\alpha = -\xi/(8\kappa)$ and $\beta = 1/(12\kappa)$. Since this expression is an odd function of ξ , we obtain (iv). Finally, we deduce (i) from (iv) by observing that, since $(3|\cdot|^{4/3}/4)^* = |\cdot|^4/4$, lemma 2.3(iii) with $\gamma = 4\kappa/3$ yields $\pi = \text{prox}_{\gamma(3|\cdot|^{4/3}/4)}\xi = \xi - \gamma \text{prox}_{(4\gamma)^{-1}|\cdot|^4}(\xi/\gamma)$, hence the result after simple algebra. \square

Example 4.5 (Huber distribution). Let $\omega \in]0, +\infty[$, $\tau \in]0, +\infty[$, and set

$$\phi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \begin{cases} \tau\xi^2, & \text{if } |\xi| \leq \omega/\sqrt{2\tau}; \\ \omega\sqrt{2\tau}|\xi| - \omega^2/2, & \text{otherwise.} \end{cases} \tag{4.7}$$

Then, for every $\xi \in \mathbb{R}$,

$$\text{prox}_\phi \xi = \begin{cases} \frac{\xi}{2\tau + 1}, & \text{if } |\xi| \leq \omega(2\tau + 1)/\sqrt{2\tau}; \\ \xi - \omega\sqrt{2\tau} \text{sign}(\xi), & \text{if } |\xi| > \omega(2\tau + 1)/\sqrt{2\tau}. \end{cases} \tag{4.8}$$

Proof. Let $\xi \in \mathbb{R}$ and set $\pi = \text{prox}_\phi \xi$. Since ϕ is even, we assume that $\xi \geq 0$ (see remark 4.1(ii)). In addition, since ϕ is differentiable, it follows from (2.7) and (4.1) that π is the unique solution in $[0, \xi]$ to $\pi + \phi'(\pi) = \xi$. First, suppose that $\pi = \omega/\sqrt{2\tau}$. Then $\phi'(\pi) = \omega\sqrt{2\tau}$ and, therefore, $\xi = \pi + \phi'(\pi) = \omega(2\tau + 1)/\sqrt{2\tau}$. Now, suppose that $\xi \leq \omega(2\tau + 1)/\sqrt{2\tau}$. Then it follows from lemma 2.6(i) that $\pi \leq \text{prox}_\phi(\omega(2\tau + 1)/\sqrt{2\tau}) = \omega/\sqrt{2\tau}$. In turn, (4.7) yields $\phi'(\pi) = 2\tau\pi$ and the identity $\xi = \pi + \phi'(\pi)$ yields $\pi = \xi/(2\tau + 1)$. Finally, if $\xi > \omega(2\tau + 1)/\sqrt{2\tau}$, then lemma 2.6(i) yields $\pi \geq \text{prox}_\phi(\omega(2\tau + 1)/\sqrt{2\tau}) = \omega/\sqrt{2\tau}$ and, in turn, $\phi'(\pi) = \omega\sqrt{2\tau}$, which allows us to conclude that $\pi = \xi - \omega\sqrt{2\tau}$. \square

Example 4.6 (maximum entropy distribution). This density is obtained by maximizing the entropy subject to the knowledge of the first, second and p th order absolute moments, where $2 \neq p \in]1, +\infty[$ [24]. Let $\omega \in]0, +\infty[$, $\tau \in [0, +\infty[$, $\kappa \in]0, +\infty[$, and set

$$\phi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \omega|\xi| + \tau|\xi|^2 + \kappa|\xi|^p. \tag{4.9}$$

Then, for every $\xi \in \mathbb{R}$,

$$\text{prox}_\phi \xi = \text{sign}(\xi)\text{prox}_{\kappa|\cdot|^p/(2\tau+1)}\left(\frac{1}{2\tau + 1} \max\{|\xi| - \omega, 0\}\right) \tag{4.10}$$

where the expression for $\text{prox}_{\kappa|\cdot|^p/(2\tau+1)}$ is supplied by example 4.4.

Proof. The function ϕ is a quadratic perturbation of the function $\varphi = \omega|\cdot| + \kappa|\cdot|^p$. Applying lemma 2.6(iii) with $\psi = \kappa|\cdot|^p$ and $\Omega = [-\omega, \omega]$, we get $(\forall \xi \in \mathbb{R}) \text{prox}_\phi \xi = \text{prox}_{\kappa|\cdot|^p}(\text{soft}_{[-\omega, \omega]}\xi) = \text{sign}(\xi)\text{prox}_{\kappa|\cdot|^p}(\max\{|\xi| - \omega, 0\})$. Hence, the result follows from lemma 2.5(i) where $\mathcal{X} = \mathbb{R}$, $\alpha = 2\tau$ and $u = 0$. \square

Example 4.7 (smoothed Laplace distribution). Let $\omega \in]0, +\infty[$ and set

$$\phi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \omega|\xi| - \ln(1 + \omega|\xi|). \quad (4.11)$$

This potential function is sometimes used as a differentiable approximation to (4.2), e.g., [29]. We have, for every $\xi \in \mathbb{R}$,

$$\text{prox}_\phi \xi = \text{sign}(\xi) \frac{\omega|\xi| - \omega^2 - 1 + \sqrt{(\omega|\xi| - \omega^2 - 1)^2 + 4\omega|\xi|}}{2\omega}. \quad (4.12)$$

Proof. According to remark 4.1(ii), since ϕ is even, we can focus on the case when $\xi \geq 0$. As ϕ achieves its infimum at 0, lemma 2.6(ii) yields $\pi = \text{prox}_\phi \xi \geq 0$. We deduce from (4.1) that π is the unique solution in $[0, +\infty[$ to the equation

$$\omega\pi^2 + (\omega^2 + 1 - \omega\xi)\pi - \xi = 0, \quad (4.13)$$

which leads to (4.12). \square

Example 4.8 (exponential distribution). Let $\omega \in]0, +\infty[$ and set

$$\phi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \begin{cases} \omega\xi, & \text{if } \xi \geq 0; \\ +\infty, & \text{if } \xi < 0. \end{cases} \quad (4.14)$$

Then, for every $\xi \in \mathbb{R}$,

$$\text{prox}_\phi \xi = \begin{cases} \xi - \omega & \text{if } \xi \geq \omega; \\ 0 & \text{if } \xi < \omega. \end{cases} \quad (4.15)$$

Proof. Set $\varphi = \iota_{[0, +\infty[}$. Then example 2.2 yields $\text{prox}_\varphi = P_{[0, +\infty[}$. In turn, since ϕ is a linear perturbation of φ , the claim results from lemma 2.5(i), where $\mathcal{X} = \mathbb{R}$, $\alpha = 0$ and $u = \omega$. \square

Example 4.9 (gamma distribution). Let $\omega \in]0, +\infty[$, $\kappa \in]0, +\infty[$, and set

$$\phi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \begin{cases} -\kappa \ln(\xi) + \omega\xi, & \text{if } \xi > 0; \\ +\infty, & \text{if } \xi \leq 0. \end{cases} \quad (4.16)$$

Then, for every $\xi \in \mathbb{R}$,

$$\text{prox}_\phi \xi = \frac{\xi - \omega + \sqrt{(\xi - \omega)^2 + 4\kappa}}{2}. \quad (4.17)$$

Proof. Set

$$\varphi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \begin{cases} -\kappa \ln(\xi), & \text{if } \xi > 0; \\ +\infty, & \text{if } \xi \leq 0. \end{cases} \quad (4.18)$$

We easily get from remark 4.1(i) and (iii) that

$$(\forall \xi \in \mathbb{R}) \quad \text{prox}_\varphi \xi = \frac{\xi + \sqrt{\xi^2 + 4\kappa}}{2}. \quad (4.19)$$

In turn, since ϕ is a linear perturbation of φ , the claim results from lemma 2.5(i), where $\mathcal{X} = \mathbb{R}$, $\alpha = 0$ and $u = \omega$. \square

Example 4.10 (chi distribution). Let $\kappa \in]0, +\infty[$ and let

$$\phi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \begin{cases} -\kappa \ln(\xi) + \xi^2/2, & \text{if } \xi > 0; \\ +\infty, & \text{if } \xi \leq 0. \end{cases} \quad (4.20)$$

Then, for every $\xi \in \mathbb{R}$,

$$\text{prox}_\phi \xi = \frac{\xi + \sqrt{\xi^2 + 8\kappa}}{4}. \tag{4.21}$$

Proof. Since ϕ is a quadratic perturbation of the function φ defined in (4.18), the claim results from lemma 2.5(i), where $\mathcal{X} = \mathbb{R}$, $\alpha = 1$ and $u = 0$. \square

Example 4.11 (uniform distribution). Let $\omega \in]0, +\infty[$ and set $\phi = \iota_{[-\omega, \omega]}$. Then it follows at once from example 2.2 that, for every $\xi \in \mathbb{R}$,

$$\text{prox}_\phi \xi = P_{[-\omega, \omega]} \xi = \begin{cases} -\omega, & \text{if } \xi < -\omega; \\ \xi, & \text{if } |\xi| \leq \omega; \\ \omega, & \text{if } \xi > \omega. \end{cases} \tag{4.22}$$

Example 4.12 (triangular distribution). Let $\underline{\omega} \in]-\infty, 0[$, let $\bar{\omega} \in]0, +\infty[$ and set

$$\phi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \begin{cases} -\ln(\xi - \underline{\omega}) + \ln(-\underline{\omega}), & \text{if } \xi \in]\underline{\omega}, 0]; \\ -\ln(\bar{\omega} - \xi) + \ln(\bar{\omega}), & \text{if } \xi \in]0, \bar{\omega}]; \\ +\infty, & \text{otherwise.} \end{cases} \tag{4.23}$$

Then, for every $\xi \in \mathbb{R}$,

$$\text{prox}_\phi \xi = \begin{cases} \frac{\xi + \underline{\omega} + \sqrt{|\xi - \underline{\omega}|^2 + 4}}{2}, & \text{if } \xi < 1/\underline{\omega}; \\ \frac{\xi + \bar{\omega} - \sqrt{|\xi - \bar{\omega}|^2 + 4}}{2}, & \text{if } \xi > 1/\bar{\omega}; \\ 0 & \text{otherwise.} \end{cases} \tag{4.24}$$

Proof. Let $\xi \in \mathbb{R}$ and set $\pi = \text{prox}_\phi \xi$. Let us first note that $\partial\phi(0) = [1/\underline{\omega}, 1/\bar{\omega}]$. Therefore, (2.5) yields

$$\pi = 0 \iff \xi \in [1/\underline{\omega}, 1/\bar{\omega}]. \tag{4.25}$$

Now consider the case when $\xi > 1/\bar{\omega}$. Since ϕ admits 0 as a minimizer, it follows from lemma 2.6(ii) and (4.25) that $\pi \in]0, \xi]$. Hence, we derive from (4.1) that π is the only solution in $]0, \xi]$ to $\pi + 1/(\bar{\omega} - \pi) = \xi$, i.e., $\pi = (\xi + \bar{\omega} - \sqrt{|\xi - \bar{\omega}|^2 + 4})/2$. Likewise, if $\xi < 1/\underline{\omega}$, it follows from lemma 2.6(ii), (4.25) and (4.1) that π is the only solution in $[\xi, 0[$ to $\pi - 1/(\pi - \underline{\omega}) = \xi$, which yields $\pi = (\xi + \underline{\omega} + \sqrt{|\xi - \underline{\omega}|^2 + 4})/2$. \square

The next example is an extension of example 4.10.

Example 4.13 (Weibull distribution). Let $\omega \in]0, +\infty[$, $\kappa \in]0, +\infty[$ and $p \in]1, +\infty[$, and set

$$\phi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \begin{cases} -\kappa \ln(\xi) + \omega\xi^p, & \text{if } \xi > 0; \\ +\infty, & \text{if } \xi \leq 0. \end{cases} \tag{4.26}$$

Then, for every $\xi \in \mathbb{R}$, $\pi = \text{prox}_\phi \xi$ is the unique strictly positive solution to

$$p\omega\pi^p + \pi^2 - \xi\pi = \kappa. \tag{4.27}$$

Proof. Since ϕ is differentiable on $]0, +\infty[$, it follows from remark 4.1(i) and (iii) that π is the unique solution in $]0, +\infty[$ to $\pi + \phi'(\pi) = \xi$ or, equivalently, to (4.27). \square

A similar proof can be used in the following two examples.

Example 4.14 (generalized inverse Gaussian distribution). Let $\omega \in]0, +\infty[$, $\kappa \in [0, +\infty[$ and $\rho \in]0, +\infty[$, and set

$$\phi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \begin{cases} -\kappa \ln(\xi) + \omega\xi + \rho/\xi, & \text{if } \xi > 0; \\ +\infty, & \text{if } \xi \leq 0. \end{cases} \tag{4.28}$$

Then, for every $\xi \in \mathbb{R}$, $\pi = \text{prox}_\phi \xi$ is the unique strictly positive solution to

$$\pi^3 + (\omega - \xi)\pi^2 - \kappa\pi = \rho. \tag{4.29}$$

Example 4.15 (Pearson type I). Let $\underline{\kappa}$ and $\bar{\kappa}$ be in $]0, +\infty[$, let $\underline{\omega}$ and $\bar{\omega}$ be reals such that $\underline{\omega} < \bar{\omega}$, and set

$$\phi : \mathbb{R} \rightarrow]-\infty, +\infty] : \xi \mapsto \begin{cases} -\underline{\kappa} \ln(\xi - \underline{\omega}) - \bar{\kappa} \ln(\bar{\omega} - \xi), & \text{if } \xi \in]\underline{\omega}, \bar{\omega}[; \\ +\infty, & \text{otherwise.} \end{cases} \tag{4.30}$$

Then, for every $\xi \in \mathbb{R}$, $\pi = \text{prox}_\phi \xi$ is the unique solution in $]\underline{\omega}, \bar{\omega}[$ to

$$\pi^3 - (\underline{\omega} + \bar{\omega} + \xi)\pi^2 + (\underline{\omega}\bar{\omega} - \underline{\kappa} - \bar{\kappa} + (\underline{\omega} + \bar{\omega})\xi)\pi = \underline{\omega}\bar{\omega}\xi - \underline{\omega}\bar{\kappa} - \bar{\omega}\underline{\kappa}. \tag{4.31}$$

Remark 4.16.

- (i) The chi-square distribution with $n > 2$ degrees of freedom is a special case of the gamma distribution (example 4.9) with $(\omega, \kappa) = (1/2, n/2 - 1)$.
- (ii) The normalized Rayleigh distribution is a special case of the chi distribution (example 4.10) with $\kappa = 1$.
- (iii) The beta distribution and the Wigner distribution are special cases of the Pearson type I distribution (example 4.15) with $(\underline{\omega}, \bar{\omega}) = (0, 1)$, and $-\underline{\omega} = \bar{\omega}$ and $\underline{\kappa} = \bar{\kappa} = 1/2$, respectively.
- (iv) The proximity operator associated with translated and/or scaled versions of the above densities can be obtained via lemma 2.5(ii)–(iii).
- (v) For log-concave densities for which the proximity operator of the potential function is difficult to express in closed form (e.g., Kumaraswamy or logarithmic distributions), one can turn to simple procedures to solve (2.5) or (4.1) numerically.

5. Algorithm

We propose the following algorithm to solve problem 3.1.

Algorithm 5.1. Fix $\mathbf{x}_0 \in \ell^2(\mathbb{K})$ and construct a sequence $(\mathbf{x}_n)_{n \in \mathbb{N}} = ((\xi_{n,k})_{k \in \mathbb{K}})_{n \in \mathbb{N}}$ by setting, for every $n \in \mathbb{N}$,

$$(\forall k \in \mathbb{K}) \quad \xi_{n+1,k} = \xi_{n,k} + \lambda_n (\text{prox}_{\gamma_n \phi_k}(\xi_{n,k} - \gamma_n(\eta_{n,k} + \beta_{n,k})) + \alpha_{n,k} - \xi_{n,k}), \tag{5.1}$$

where $\lambda_n \in]0, 1]$, $\gamma_n \in]0, +\infty[$, $\{\alpha_{n,k}\}_{k \in \mathbb{K}} \subset \mathbb{R}$, $(\eta_{n,k})_{k \in \mathbb{K}} = F(\nabla \Psi(F^* \mathbf{x}_n))$, and $(\beta_{n,k})_{k \in \mathbb{K}} = Fb_n$, where $b_n \in \mathcal{H}$.

The chief advantage of this algorithm is to be fully split in the sense that the functions $(\phi_k)_{k \in \mathbb{K}}$ and Ψ appearing in (3.2) are used separately. First, the current iterate \mathbf{x}_n is transformed into a point in $F^* \mathbf{x}_n$ in \mathcal{H} , and the gradient of Ψ is evaluated at this point to within some tolerance b_n . Next, we obtain the sequence $(\eta_{n,k})_{k \in \mathbb{K}} = F(\nabla \Psi(F^* \mathbf{x}_n))$ to within some tolerance $(\beta_{n,k})_{k \in \mathbb{K}} = Fb_n$. Then one chooses $\gamma_n > 0$, and, for every $k \in \mathbb{K}$, applies the

operator $\text{prox}_{\gamma_n \phi_k}$ to $\xi_{n,k} - \gamma_n(\eta_{n,k} + \beta_{n,k})$. An error $\alpha_{n,k}$ is tolerated in this computation. Finally, the k th component $\xi_{n+1,k}$ of \mathbf{x}_{n+1} is obtained by applying a relaxation of parameter λ_n to this inexact proximal step. Let us note that the computation of the proximal steps can be performed in parallel.

To study the asymptotic behaviour of the sequences generated by algorithm 5.1, we require the following set of assumptions.

Assumption 5.2. *In addition to the standing assumptions of problem 3.1, the following hold.*

- (i) *Problem 3.1 admits a solution.*
- (ii) $\inf_{n \in \mathbb{N}} \lambda_n > 0$.
- (iii) $\inf_{n \in \mathbb{N}} \gamma_n > 0$ and $\sup_{n \in \mathbb{N}} \gamma_n < 2/\beta$, where β is a Lipschitz constant of $F \circ \nabla \Psi \circ F^*$.
- (iv) $\sum_{n \in \mathbb{N}} \sqrt{\sum_{k \in \mathbb{K}} |\alpha_{n,k}|^2} < +\infty$ and $\sum_{n \in \mathbb{N}} \|b_n\| < +\infty$.

Remark 5.3. As regards assumption 5.2(i), sufficient conditions can be found in proposition 3.3. Let us now turn to the parameter β in assumption 5.2(iii), which determines the range of the step sizes $(\gamma_n)_{n \in \mathbb{N}}$. It follows from the assumptions of problem 3.1 and (1.1) that, for every \mathbf{x} and \mathbf{y} in $\ell^2(\mathbb{K})$,

$$\begin{aligned} \|F(\nabla \Psi(F^* \mathbf{x})) - F(\nabla \Psi(F^* \mathbf{y}))\| &\leq \|F\| \|\nabla \Psi(F^* \mathbf{x}) - \nabla \Psi(F^* \mathbf{y})\| \\ &\leq \tau \|F\| \|F^* \mathbf{x} - F^* \mathbf{y}\| \\ &\leq \tau \|F\|^2 \|\mathbf{x} - \mathbf{y}\| \\ &\leq \tau \nu \|\mathbf{x} - \mathbf{y}\|. \end{aligned} \tag{5.2}$$

Thus, the value $\beta = \tau \nu$ can be used in general. In some cases, however, a sharper bound can be obtained, which results in a wider range for the step sizes $(\gamma_n)_{n \in \mathbb{N}}$. For example, in the problem considered in section 3.3.1, if the norm of $R = \sum_{i=1}^q \alpha_i F T_i^* T_i F^*$ can be evaluated, it follows from (3.13) and the nonexpansivity of the operators $(\text{Id} - P_{S_i})_{1 \leq i \leq m}$ that one can take

$$\beta = \|R\| + \nu \sum_{i=1}^m \vartheta_i. \tag{5.3}$$

Theorem 5.4. *Let $(\mathbf{x}_n)_{n \in \mathbb{N}}$ be an arbitrary sequence generated by algorithm 5.1 under assumption 5.2. Then $(\mathbf{x}_n)_{n \in \mathbb{N}}$ converges weakly to a solution to problem 3.1.*

Proof. Set $\mathcal{X} = \ell^2(\mathbb{K})$, $f_1 = \sum_{k \in \mathbb{K}} \phi_k(\cdot | \mathbf{o}_k)$ and $f_2 = \Psi \circ F^*$, where $(\mathbf{o}_k)_{k \in \mathbb{K}}$ denotes the canonical orthonormal basis of $\ell^2(\mathbb{K})$. Then $\nabla f_2 = F \circ \nabla \Psi \circ F^*$ is β -Lipschitz continuous (see assumption 5.2(iii)) and, as seen in the proof of proposition 3.4, (3.2) conforms to the format of problem 2.7. Furthermore, it follows from proposition 2.10 that we can rewrite (5.1) as

$$\begin{aligned} \mathbf{x}_{n+1} &= \mathbf{x}_n + \lambda_n \left(\sum_{k \in \mathbb{K}} (\text{prox}_{\gamma_n \phi_k}(\mathbf{x}_n - \gamma_n F(\nabla \Psi(F^* \mathbf{x}_n)) + b_n) | \mathbf{o}_k) + \alpha_{n,k} \mathbf{o}_k - \mathbf{x}_n \right) \\ &= \mathbf{x}_n + \lambda_n (\text{prox}_{\gamma_n f_1}(\mathbf{x}_n - \gamma_n (\nabla f_2(\mathbf{x}_n) + \mathbf{b}_n)) + \mathbf{a}_n - \mathbf{x}_n), \end{aligned} \tag{5.4}$$

where $\mathbf{a}_n = (\alpha_{n,k})_{k \in \mathbb{K}}$ and $\mathbf{b}_n = F b_n$. Since assumptions 5.2(iv) and (1.1) imply that $\sum_{n \in \mathbb{N}} \|\mathbf{a}_n\| < +\infty$ and $\sum_{n \in \mathbb{N}} \|\mathbf{b}_n\| \leq \sqrt{\nu} \sum_{n \in \mathbb{N}} \|b_n\| < +\infty$, the claim therefore follows from theorem 2.9. \square

Let $(\mathbf{x}_n)_{n \in \mathbb{N}}$ be a sequence generated by algorithm 5.1 under assumption 5.2 and set $(\forall n \in \mathbb{N}) \mathbf{x}_n = F^* \mathbf{x}_n$. On the one hand, theorem 5.4 asserts that $(\mathbf{x}_n)_{n \in \mathbb{N}}$ converges weakly to

a solution x to problem 3.1. On the other hand, since F^* is linear and bounded, it is weakly continuous and, therefore, $(x_n)_{n \in \mathbb{N}}$ converges weakly to F^*x . However, it is not possible to express (5.1) as an iteration in terms of the sequence $(x_n)_{n \in \mathbb{N}}$ in \mathcal{H} in general. The following corollary addresses the case when F is surjective, which does lead to an algorithm in \mathcal{H} .

Corollary 5.5. *Suppose that $(e_k)_{k \in \mathbb{K}}$ is a Riesz basis with companion biorthogonal basis $(\check{e}_k)_{k \in \mathbb{K}}$. Fix $x_0 \in \mathcal{H}$ and, for every $n \in \mathbb{N}$, set*

$$x_{n+1} = x_n + \lambda_n \left(\sum_{k \in \mathbb{K}} (\text{prox}_{\gamma_n \phi_k} (\langle x_n | \check{e}_k \rangle - \gamma_n \langle \nabla \Psi(x_n) + b_n | e_k \rangle) + \alpha_{n,k}) e_k - x_n \right), \quad (5.5)$$

where $\lambda_n \in]0, 1]$, $\gamma_n \in]0, +\infty[$, $\{\alpha_{n,k}\}_{k \in \mathbb{K}} \subset \mathbb{R}$ and $b_n \in \mathcal{H}$. Suppose that assumption 5.2 is in force. Then $(x_n)_{n \in \mathbb{N}}$ converges weakly to a point $x \in \mathcal{H}$ and $(\langle x | \check{e}_k \rangle)_{k \in \mathbb{K}}$ is a solution to problem 3.1.

Proof. Set $(\forall n \in \mathbb{N}) (\forall k \in \mathbb{K}) \xi_{n,k} = \langle x_n | \check{e}_k \rangle$, $\eta_{n,k} = \langle \nabla \Psi(x_n) | e_k \rangle$ and $\beta_{n,k} = \langle b_n | e_k \rangle$. Then, for every $n \in \mathbb{N}$, it follows from (1.4) that $x_n = F^*(\xi_{n,k})_{k \in \mathbb{K}}$ and, in turn, that

$$(\eta_{n,k})_{k \in \mathbb{K}} = F(\nabla \Psi(F^*x_n)), \quad \text{where } x_n = (\xi_{n,k})_{k \in \mathbb{K}}. \quad (5.6)$$

Furthermore, for every $n \in \mathbb{N}$, it follows from (5.5) and the biorthogonality of $(e_k)_{k \in \mathbb{K}}$ and $(\check{e}_k)_{k \in \mathbb{K}}$ that

$$\begin{aligned} (\forall k \in \mathbb{K}) \quad \xi_{n+1,k} &= \langle x_{n+1} | \check{e}_k \rangle \\ &= \langle x_n | \check{e}_k \rangle + \lambda_n (\text{prox}_{\gamma_n \phi_k} (\langle x_n | \check{e}_k \rangle - \gamma_n \langle \nabla \Psi(x_n) + b_n | e_k \rangle) \\ &\quad + \alpha_{n,k} - \langle x_n | \check{e}_k \rangle) \\ &= \xi_{n,k} + \lambda_n (\text{prox}_{\gamma_n \phi_k} (\xi_{n,k} - \gamma_n (\eta_{n,k} + \beta_{n,k})) + \alpha_{n,k} - \xi_{n,k}). \end{aligned} \quad (5.7)$$

Since theorem 5.4 states that $(x_n)_{n \in \mathbb{N}}$ converges weakly to a solution x to problem 3.1, $(x_n)_{n \in \mathbb{N}} = (F^*x_n)_{n \in \mathbb{N}}$ converges weakly to $x = F^*x$. Consequently, (1.4) asserts that we can write $x = (\langle x | \check{e}_k \rangle)_{k \in \mathbb{K}}$. \square

Remark 5.6. Suppose that $\mathbb{K} = \mathbb{N}$ and that $(e_k)_{k \in \mathbb{K}}$ is an orthonormal basis of \mathcal{H} . Then (5.5) reduces to

$$x_{n+1} = x_n + \lambda_n \left(\sum_{k \in \mathbb{K}} (\text{prox}_{\gamma_n \phi_k} (\langle x_n - \gamma_n (\nabla \Psi(x_n) + b_n) | e_k \rangle) + \alpha_{n,k}) e_k - x_n \right). \quad (5.8)$$

In this particular setting, some results related to corollary 5.5 are the following.

- (i) Suppose that $\Psi : x \mapsto \|Tx - z\|^2/2$, where T is a nonzero bounded linear operator from \mathcal{H} to a real Hilbert space \mathcal{G} and $z \in \mathcal{G}$. Suppose that, in addition, $(\forall k \in \mathbb{K}) \phi_k \geq \phi_k(0) = 0$. Then the convergence of (5.8) is discussed in [15, corollary 5.16].
- (ii) Suppose that $(\Omega_k)_{k \in \mathbb{K}}$ are closed intervals of \mathbb{R} such that $0 \in \text{int} \bigcap_{k \in \mathbb{K}} \Omega_k$ and that

$$(\forall k \in \mathbb{K}) \quad \phi_k = \psi_k + \sigma_{\Omega_k}, \quad (5.9)$$

where $\psi_k \in \Gamma_0(\mathbb{R})$ is differentiable at 0 and $\psi_k \geq \psi_k(0) = 0$. Then (5.8) is the thresholding algorithm proposed and analysed in [14], namely

$$x_{n+1} = x_n + \lambda_n \left(\sum_{k \in \mathbb{K}} (\text{prox}_{\gamma_n \psi_k} (\text{soft}_{\gamma_n \Omega_k} \langle x_n - \gamma_n (\nabla \Psi(x_n) + b_n) | e_k \rangle) + \alpha_{n,k}) e_k - x_n \right), \quad (5.10)$$

where $\text{soft}_{\gamma_n \Omega_k}$ is defined in (2.8).

- (iii) Suppose that the assumptions of both (i) and (ii) hold and that, in addition, we set $\lambda_n \equiv 1$, $\|T\| < 1$, $\gamma_n \equiv 1$, $\alpha_{n,k} \equiv 0$, $b_n \equiv 0$, and $(\forall k \in \mathbb{K}) \psi_k = 0$ and $\Omega_k = [-\omega_k, \omega_k]$. Then (5.10) becomes

$$x_{n+1} = \sum_{k \in \mathbb{K}} (\text{soft}_{[-\omega_k, \omega_k]}(x_n + T^*(z - Tx_n) \mid e_k)) e_k. \tag{5.11}$$

Algorithm 5.1 can be regarded as a descendant of this original method, which is investigated in [17, 18].

6. Numerical results

The proposed framework is applicable to a wide array of variational formulations for inverse problems over frames. We provide a couple of examples to illustrate its applicability in wavelet-based image restoration in the Euclidean space $\mathcal{H} = \mathbb{R}^{512 \times 512}$. The choice of the potential functions $(\phi_k)_{k \in \mathbb{K}}$ in problem 3.1 is guided by the observation that regular images typically possess sparse wavelet representations and that the resulting wavelet coefficients often have even probability density functions [26]. Among the candidate potential functions investigated in section 4, those of example 4.6 appear to be the most appropriate for modelling wavelet coefficients on two counts. First, they provide flexible models of even potentials. Second, as shown in lemma 2.6(iii), their proximity operators are thresholders and they therefore promote sparsity. More precisely, we employ potential functions of the form $\phi_k = \omega_k |\cdot| + \tau_k |\cdot|^2 + \kappa_k |\cdot|^{p_k}$, where $p_k \in \{4/3, 3/2, 3, 4\}$ and $\{\omega_k, \tau_k, \kappa_k\} \subset]0, +\infty[$. Note that prox_{ϕ_k} can be obtained explicitly via (4.10) and examples 4.4(i)–(iv). In addition, it follows from proposition 3.3(iii)(b) that, with such potential functions, problem 3.1 does admit a solution. The values of the parameters $\omega_k, \tau_k, \kappa_k$ and p_k are chosen for each wavelet subband via a maximum likelihood approach. The first example uses a biorthogonal wavelet basis and the second one uses an M -band dual-tree wavelet frame. Let us emphasize that such decompositions cannot be dealt with using the methods developed in [14], which are limited to orthonormal basis representations. Algorithm 5.1 is implemented with $\lambda_n \equiv 1$ and large step sizes (i.e., γ_n close to $2/\beta$) since such values have been observed to provide a good speed of convergence in our experiments.

6.1. Example 1

We provide a multiview restoration example in a biorthogonal wavelet basis. The original image \bar{x} is the standard test image displayed in figure 2 (top left). Two observations (see figure 2, top right and bottom left) conforming to the model (3.12) are available. In our experiment, $\mathcal{G}_1 = \mathcal{G}_2 = \mathcal{H}$ and v_1 and v_2 are realizations of two independent zero-mean Gaussian white noise processes. Moreover, the operator T_1 models a motion blur in the diagonal direction and satisfies $\|T_1\| = 1$, whereas $T_2 = \text{Id}/2$. The blurred image-to-noise ratio is higher for the first observation (22.79 dB versus 15.18 dB) and so is the relative error (18.53 dB versus 5.891 dB) (the decibel value of the relative error between an image z and \bar{x} is $20 \log_{10}(\|\bar{x}\|/\|z - \bar{x}\|)$). The function Ψ in problem 3.1 is given by (3.13), where $\alpha_1 = 4.00 \times 10^{-2}$ and $\alpha_2 = 6.94 \times 10^{-3}$ are the inverses of the variances of the noise corrupting each observation. In addition, we set $m = 1$, $\vartheta_1 = 10^{-2}$ and $S_1 = [0, 255]^{512 \times 512}$ to enforce the known range of the pixel values. A discrete biorthogonal spline 9–7 decomposition [2] is used over three resolution levels. Algorithm 5.1 is used to solve problem 3.1. By numerically evaluating $\|R\|$ in (5.3), we obtain $\beta = 0.230$ and the step sizes are chosen to be $\gamma_n \equiv 1.99/\beta = 8.66$. The resulting restored image, shown in figure 2 (bottom right), yields a relative error of 23.84 dB.




Figure 2. Example 1—original image (top left), first observation (top right), second observation (bottom left) and image restored with 200 iterations of algorithm 5.1 (bottom right).

6.2. Example 2

The original SPOT5 satellite image \bar{x} is shown in figure 3 (top) and the degraded image z in $\mathcal{G} = \mathcal{H}$ is shown in figure 3 (centre). The degradation model is given by (3.16), where T is a 7×7 uniform blur with $\|T\| = 1$, and where v is a realization of a zero-mean Gaussian white noise process. The blurred image-to-noise ratio is 28.08 dB and the relative error is 12.49 dB.

In this example, we perform a restoration in a discrete two-dimensional version of an M -band dual-tree wavelet frame [7]. This decomposition has a redundancy factor of 2 (i.e., with the notation of section 3.3.2, $K/N = 2$). In our experiments, decompositions over two resolution levels are performed with $M = 4$ using the filter bank proposed in [1]. The function Ψ in problem 3.1 is given by (3.22), where f_V is the probability density function of the Gaussian noise. A solution is obtained via algorithm 5.1. For the representation under consideration, we derive from (5.3) that $\beta = 2$ and we set $\gamma_n \equiv 0.995$. The restored image,




Figure 3. Example 2—original image (top); degraded image (centre); image restored in a dual-tree wavelet frame with 100 iterations of algorithm 5.1 (bottom).




Figure 4. Example 2—zoom on a 100×100 portion of the SPOT5 satellite image. Original image (top), degraded image (centre), image restored in a dual-tree wavelet frame with 100 iterations of algorithm 5.1 (bottom).

shown in figure 3 (bottom), yields a relative error of 15.68 dB, i.e., a significant improvement of over 3 dB in terms of signal-to-noise ratio. A more precise inspection of the magnified areas displayed in figure 4 shows that the proposed method makes it possible to recover sharp edges while removing noise in uniform areas. This behaviour in terms of edge recovery may be attributed to the choice of the M -band dual-tree wavelet decomposition, which is known to provide a good representation of directional features such as edges [7].

References

- [1] Alkin O and Caglar H 1995 Design of efficient M -band coders with linear-phase and perfect-reconstruction properties *IEEE Trans. Signal Process.* **43** 1579–90
- [2] Antonini M, Barlaud M, Mathieu P and Daubechies I 1992 Image coding using wavelet transform *IEEE Trans. Image Process.* **1** 205–20
- [3] Bioucas-Dias J M 2006 Bayesian wavelet-based image deconvolution: a GEM algorithm exploiting a class of heavy-tailed priors *IEEE Trans. Image Process.* **15** 937–51
- [4] Bouman C and Sauer K 1993 A generalized Gaussian image model for edge-preserving MAP estimation *IEEE Trans. Image Process.* **2** 296–310
- [5] Byrne C L 2004 A unified treatment of some iterative algorithms in signal processing and image reconstruction *Inverse Problems* **20** 103–20
- [6] Candès E J and Donoho D L 2002 Recovering edges in ill-posed inverse problems: optimality of curvelet frames *Ann. Stat.* **30** 784–842
- [7] Chaux C, Duval L and Pesquet J-C 2006 Image analysis using a dual-tree M -band wavelet transform *IEEE Trans. Image Process.* **15** 2397–412
- [8] Cohen A 2003 *Numerical Analysis of Wavelet Methods* (New York: Elsevier)
- [9] Cohen A, Daubechies I and Feauveau J-C 1992 Biorthogonal bases of compactly supported wavelets *Commun. Pure Appl. Math.* **45** 485–560
- [10] Combettes P L 1993 The foundations of set theoretic estimation *Proc. IEEE* **81** 182–208
- [11] Combettes P L 1994 Inconsistent signal feasibility problems: least-squares solutions in a product space *IEEE Trans. Signal Process.* **42** 2955–66
- [12] Combettes P L 2003 A block-iterative surrogate constraint splitting method for quadratic signal recovery *IEEE Trans. Signal Process.* **51** 1771–82
- [13] Combettes P L 2004 Solving monotone inclusions via compositions of nonexpansive averaged operators *Optimization* **53** 475–504
- [14] Combettes P L and Pesquet J-C 2007 Proximal thresholding algorithm for minimization over orthonormal bases *SIAM J. Optim.* at press
- [15] Combettes P L and Wajs V R 2005 Signal recovery by proximal forward–backward splitting *Multiscale Model. Simul.* **4** 1168–200
- [16] Daubechies I 1992 *Ten Lectures on Wavelets* (Philadelphia, PA: SIAM)
- [17] Daubechies I, Defrise M and De Mol C 2004 An iterative thresholding algorithm for linear inverse problems with a sparsity constraint *Commun. Pure Appl. Math.* **57** 1413–57
- [18] De Mol C and Defrise M 2002 A note on wavelet-based inversion algorithms *Contemp. Math.* **313** 85–96
- [19] Devroye L 1986 *Non-Uniform Random Variate Generation* (New York: Springer)
- [20] Do M N and Vetterli M 2005 The contourlet transform: an efficient directional multiresolution image representation *IEEE Trans. Image Process.* **14** 2091–106
- [21] Elad M and Feuer A 1997 Restoration of a single superresolution image from several blurred, noisy, and undersampled measured images *IEEE Trans. Image Process.* **6** 1646–58
- [22] Escoda O D, Granai L and Vandergheynst P 2006 On the use of *a priori* information for sparse signal approximations *IEEE Trans. Signal Process.* **54** 3468–82
- [23] Han D and Larson D R 2000 *Frames, Bases, and Group Representations* (*Mem. Am. Math. Soc.* vol 147)
- [24] Kapur J N and Keshavan H K 1992 *Entropy Optimization Principles with Applications* (Boston, MA: Academic)
- [25] Le Pennec E and Mallat S G 2005 Sparse geometric image representations with bandelets *IEEE Trans. Image Process.* **14** 423–38
- [26] Mallat S G 1999 *A Wavelet Tour of Signal Processing* 2nd edn (New York: Academic)
- [27] Moreau J-J 1962 Fonctions convexes duales et points proximaux dans un espace hilbertien *C. R. Acad. Sci. Paris A* **255** 2897–9
- [28] Moreau J-J 1965 Proximité et dualité dans un espace hilbertien *Bull. Soc. Math. France* **93** 273–99

-
- [29] Nikolova M and Ng M K 2005 Analysis of half-quadratic minimization methods for signal and image recovery *SIAM J. Sci. Comput.* **27** 937–66
 - [30] Selesnick I W, Baraniuk R G and Kingsbury N C 2005 The dual-tree complex wavelet transform *IEEE Signal Process. Mag.* **22** 123–51
 - [31] Stark H (ed) 1987 *Image Recovery: Theory and Application* (San Diego, CA: Academic)
 - [32] Thompson A M and Kay J 1993 On some Bayesian choices of regularization parameter in image restoration *Inverse Problems* **9** 749–61
 - [33] Tolimieri R and An M 1998 *Time-Frequency Representations* (Boston, MA: Birkhäuser)
 - [34] Tropp J A 2006 Just relax: convex programming methods for identifying sparse signals in noise *IEEE Trans. Inform. Theory* **52** 1030–51
 - [35] Van Trees H L 1968 *Detection, Estimation, and Modulation Theory—Part I* (New York: Wiley)
 - [36] Zălinescu C 2002 *Convex Analysis in General Vector Spaces* (River Edge, NJ: World Scientific)