

# A New Approach for Mutual Information Analysis of Large Dimensional Multi-Antenna Channels

Walid Hachem, *Member, IEEE*, Oleksiy Khorunzhiy, Philippe Loubaton, *Fellow, IEEE*, Jamal Najim, and Leonid Pastur

**Abstract**—This paper addresses the behavior of the mutual information of correlated multiple-input multiple-output (MIMO) Rayleigh channels when the numbers of transmit and receive antennas converge to  $+\infty$  at the same rate. Using a new and simple approach based on Poincaré–Nash inequality and on an integration by parts formula, it is rigorously established that the mutual information when properly centered and rescaled converges to a standard Gaussian random variable. Simple expressions for the centering and scaling parameters are provided. These results confirm previous evaluations based on the powerful but nonrigorous replica method. It is believed that the tools that are used in this paper are simple, robust, and of interest for the communications engineering community.

**Index Terms**—Central limit theorem (CLT), correlated multiple-input multiple-output (MIMO) channels, large random matrix theory, mutual information, Poincaré–Nash inequality.

## I. INTRODUCTION

IT is widely known that high spectral efficiencies are attained when multiple antennas are used at both the transmitter and the receiver of a wireless communication system. Indeed, consider the classical transmission model  $\mathbf{y} = \mathbf{G}\mathbf{x} + \mathbf{z}$ , where  $\mathbf{y}$  is the received signal,  $\mathbf{x}$  is the vector of transmitted symbols,  $\mathbf{z}$  is a complex white Gaussian noise, and  $\mathbf{G}$  is the  $N \times n$  multiple-input multiple-output (MIMO) channel matrix with  $N$  antennas at the receiver's site and  $n$  antennas at the transmitter's. Due to the mobility and to the presence of a large number of reflected and scattered signal paths, the elements of the channel matrix  $\mathbf{G}$  are often modeled as random variables. Assuming a random model for this matrix, Telatar [1] and Foschini [2] realized in the mid-1990s that Shannon's mutual information of such channels increases at the rate of  $\min(N, n)$  for a fixed transmission power [1]. The authors of [1] and [2] assumed

that the elements of the channel matrix  $\mathbf{G}$  are centered, independent and identically distributed (i.i.d.) elements. In this context, a well-known result in Random Matrix Theory (RMT) [3] states that the eigenvalue distribution of the Gram matrix  $\mathbf{G}\mathbf{G}^*$  where  $\mathbf{G}^*$  is the Hermitian adjoint of  $\mathbf{G}$  converges to a deterministic probability distribution as  $n$  goes to infinity and  $N/n$  converges to a constant  $c > 0$ . Denote by  $I(\rho) = \log \det \left( \frac{\rho}{n} \mathbf{G}\mathbf{G}^* + \mathbf{I}_N \right)$  the mutual information of channel  $\mathbf{G}$  for a signal-to-noise ratio at a receiver antenna equal to  $\rho/n$ . One consequence of [3] is that the mutual information per transmit antenna  $I(\rho)/n$ , being an integral of a log function with respect to the empirical eigenvalue distribution of  $\mathbf{G}\mathbf{G}^*$ , converges to a constant. This fact already observed in [1] sustains the assertion of a linear increase of mutual information with the number of antennas. In addition, this convergence proves to be sufficiently fast. As a matter of fact, the asymptotic results predicted by the RMT remain relevant for systems with a moderate number of antennas.

The next step was to apply this theory to channel models that include a correlation between paths (or entries of  $\mathbf{G}$ ). One of the main purposes of this generalization is to better understand the impact of these correlations on Shannon's mutual information. Let us cite in this context the contributions [4]–[8], all devoted to the study of the mutual information in the case where the elements of channel's matrix are centered and correlated random variables. In [9], a deterministic equivalent is computed under broad conditions for the mutual information based on Rice channels modeled by noncentered matrices with independent but not identically distributed random variables. The link between matrices with correlated entries and matrices with independent entries and a variance profile is studied in [10].

One of the most popular correlated channel models used for these mutual information evaluations is the so-called Kronecker model  $\mathbf{G} = \mathbf{\Psi}\mathbf{W}\tilde{\mathbf{\Psi}}$  where  $\mathbf{W}$  is a  $N \times n$  matrix with Gaussian centered i.i.d. entries, and  $\mathbf{\Psi}$  and  $\tilde{\mathbf{\Psi}}$  are  $N \times N$  and  $n \times n$  matrices that capture the path correlations at the receiver and at the transmitter sides, respectively, [11], [12]. This model has been studied by Chuah *et al.* in [5]. With some assumptions on matrices  $\mathbf{\Psi}$  and  $\tilde{\mathbf{\Psi}}$ , these authors showed that  $I(\rho)/n$  converges to a deterministic quantity defined as the fixed point of an integral equation. Later on, Tulino *et al.* [8] obtained the limit of  $I(\rho)/n$  for a correlation model more general than the Kronecker model. Both these works rely on a result of Girko describing the eigenvalue distribution of the Gram matrix associated with a matrix with independent but not necessarily identically distributed entries, a close model as we shall see in a moment.

In [7], Moustakas *et al.* studied the mutual information for the Kronecker model by using the so-called replica method. They

Manuscript received December 15, 2006; revised November 20, 2007. Published August 27, 2008 (projected). This work was partially supported by the "Fonds National de la Science" via the ACI program "Nouvelles Interfaces des Mathématiques," project MALCOM number 205.

W. Hachem and J. Najim are with the CNRS (UMR 5141) and Telecom ParisTech (ENST), 75634 Paris Cedex 13, Paris, France (e-mail: najim@enst.fr; walid.hachem@enst.fr).

O. Khorunzhiy is with the Equipe "Probabilités-Statistiques" Université de Versailles—Saint-Quentin, Bureau 2306, Bâtiment Fermat, 78035 Versailles, France (e-mail: khorunjy@math.uvsq.fr).

P. Loubaton is with the IGM LabInfo, (CNRS/UMR 8049), Université Paris-Est, Equipe Signal et Communications, Institut Gaspard Monge/Electronique, 77454 Marne-la-Vallée Cedex 2, France (e-mail: loubaton@univ-mlv.fr).

L. Pastur is with the Kharkiv National University, Institute for Low Temperature Physics—Kharkiv, Kharkov 61103, Ukraine (e-mail: lpastur@flint.ilt.kharkov.ua).

Communicated by A. J. Grant, Associate Editor for Communications.  
Digital Object Identifier 10.1109/TIT.2008.928229

found an approximation  $V(\rho)$  of  $\mathbb{E}[I(\rho)]$  accurate to the order  $1/n$  in the large  $n$  regime. Using this same method, they also showed that the variance of  $I(\rho) - V(\rho)$  is of order one and were able to derive this variance for large  $n$ .

Although the replica technique is powerful and has a wide range of applications, the rigorous justification of some of its parts remains to be done. In this paper, we propose a new method to study the convergence of  $\mathbb{E}I(\rho)$  and the fluctuations of  $I(\rho)$ . Beside recovering the results in [7] and especially the strikingly simple form of the variance, we establish the Central Limit Theorem (CLT) for  $I(\rho) - V(\rho)$  (for a related CLT in a non-Gaussian context, see [13]). The practical interest of such a result is of importance since the CLT leads to an evaluation of the outage probability, i.e., the probability that  $I(\rho)$  lies beneath a given threshold, by means of the Gaussian approximation. Many other works have been devoted to CLT for random matrices. Close to our present article are [14]–[16].

In this article, we also would like to advocate the method used to establish both the approximation of  $I(\rho)$  in the large  $n$  regime and the CLT. Due to the Gaussian nature of the entries of Matrix  $\mathbf{G}$ , two simple ingredients are available. The first one is an Integration by parts formula (17) that provides an expression for the expectation of certain functionals of Gaussian vectors. This formula has been widely used in RMT [17]–[19]. The second ingredient is Poincaré–Nash inequality (18) that bounds the variance of functionals of Gaussian vectors. Although well known [20], [21], its application to RMT is fairly recent [19], [22] (see also [23] and [24] where general concentration inequalities are derived for functions of random matrices). This inequality enables us to control the decrease rate of the approximation errors such as the order  $1/n$  error  $\mathbb{E}[I(\rho)] - V(\rho)$  (note that the Gaussian structure enters in two places: First the reduction to matrices with independent entries and varying variance and then integration by part and Poincaré–Nash bounds for the variance of relevant spectral characteristics<sup>1</sup>). We believe that these tools of rigorous and explicit analysis might be of great interest for the communications engineering community (see, for instance, the estimates obtained in [25] in the context of Ricean MIMO channels).

The paper is organized as follows. In Section II, we introduce the main notations; we also state the two main results of the article. In Section III, we recall general matrix results and the two aforementioned Gaussian tools. Section IV is devoted to the proof of the first order result, that is the approximation of  $\mathbb{E}[I(\rho)]$ . The CLT, also referred to as the second order result, is established in Section V. Proof details are postponed to the Appendix.

## II. NOTATIONS AND STATEMENT OF THE MAIN RESULTS

### A. From a Kronecker Model to a Separable Variance Model

Consider a MIMO system represented by a  $N \times n$  matrix  $\mathbf{G}$  where  $n$  is the number of antennas at the transmitter and  $N$

<sup>1</sup>It is interesting to note that once the first reduction has been made, others techniques are available without assuming the Gaussian character—see for instance [19].

is the number of antennas at the receiver and where  $N(n)$  is a sequence of integers such that

$$0 < \ell^- = \liminf_{n \rightarrow \infty} \frac{N(n)}{n} \leq \ell^+ = \limsup_{n \rightarrow \infty} \frac{N(n)}{n} < \infty \quad (1)$$

a condition we shall refer to by writing  $n, N \rightarrow \infty$ . Assuming the transmitted signal is a Gaussian signal with a covariance matrix equal to  $\frac{1}{n}\mathbf{I}_n$  (and thus, a total power equal to one), Shannon's mutual information of this channel is  $I_n(\rho) = \log \det \left( \frac{\rho}{n} \mathbf{G}_n \mathbf{G}_n^* + \mathbf{I}_N \right)$ , where  $\rho > 0$  is the inverse of the additive white Gaussian noise variance at each receive antenna. The general problem we address in this paper concerns the behavior of the mutual information for large values of  $N$  and  $n$  in the case where the channel matrix  $\mathbf{G}_n$ , assumed to be random, is described by the Kronecker model  $\mathbf{G}_n = \mathbf{\Psi}_n \mathbf{W}_n \tilde{\mathbf{V}}_n$ . In this model,  $\mathbf{\Psi}_n$  and  $\tilde{\mathbf{V}}_n$  are, respectively,  $N \times N$  and  $n \times n$  deterministic matrices and  $\mathbf{W}_n$  is random with independent entries distributed according to the complex circular Gaussian law with mean zero and variance one  $\mathcal{CN}(0, 1)$ .

It is well known that this model can be replaced with a simpler Kronecker model involving a matrix with Gaussian independent (but not necessarily identically distributed) entries. Indeed, let  $\mathbf{\Psi}_n = \mathbf{U}_n \mathbf{D}_n^{\frac{1}{2}} \mathbf{V}_n^*$  (resp.,  $\tilde{\mathbf{V}}_n = \tilde{\mathbf{U}}_n \tilde{\mathbf{D}}_n^{\frac{1}{2}} \tilde{\mathbf{V}}_n^*$ ) be a Singular Value Decomposition (SVD) of  $\mathbf{\Psi}_n$  (resp.,  $\tilde{\mathbf{V}}_n$ ), where  $\mathbf{D}_n$  (resp.,  $\tilde{\mathbf{D}}_n$ ) is the diagonal matrix of eigenvalues of  $\mathbf{\Psi}_n \mathbf{\Psi}_n^*$  (resp.,  $\tilde{\mathbf{V}}_n \tilde{\mathbf{V}}_n^*$ ), then  $I_n(\rho)$  writes

$$I_n(\rho) = \log \det \left( \frac{\rho}{n} \mathbf{Y}_n \mathbf{Y}_n^* + \mathbf{I}_N \right)$$

where  $\mathbf{Y}_n = \mathbf{D}_n^{\frac{1}{2}} \mathbf{X}_n \tilde{\mathbf{D}}_n^{\frac{1}{2}}$  is a  $N \times n$  matrix,  $\mathbf{D}_n$  and  $\tilde{\mathbf{D}}_n$  are, respectively,  $N \times N$  and  $n \times n$  diagonal matrices, i.e.

$$\mathbf{D}_n = \text{diag} \left( d_i^{(n)}, 1 \leq i \leq N \right)$$

and

$$\tilde{\mathbf{D}}_n = \text{diag} \left( \tilde{d}_j^{(n)}, 1 \leq j \leq n \right)$$

and  $\mathbf{X}_n = \mathbf{V}_n^* \mathbf{W}_n \tilde{\mathbf{U}}_n$  has i.i.d. entries with distribution  $\mathcal{CN}(0, 1)$  since  $\mathbf{V}_n$  and  $\tilde{\mathbf{U}}_n$  are deterministic unitary matrices. Since every individual entry of  $\mathbf{Y}_n$  has the form  $Y_{ij}^{(n)} = \sqrt{d_i^{(n)} \tilde{d}_j^{(n)}} X_{ij}$ , we call  $\mathbf{Y}_n$  a random matrix with a separable variance profile.

### B. Assumptions and Notations

The centered random variable  $X - \mathbb{E}[X]$  will be denoted by  $\hat{X}$ . Element  $(i, j)$  of a matrix  $\mathbf{A}$  will be either denoted  $[\mathbf{A}]_{ij}$  or  $A_{ij}$ . Element  $i$  of vector  $\mathbf{a}$  will be denoted  $a_i$  or  $[\mathbf{a}]_i$ . Column  $j$  of matrix  $\mathbf{A}$  will be denoted  $\mathbf{a}_j$ . The transpose, the Hermitian adjoint (conjugate transpose) of  $\mathbf{A}$ , and the matrix obtained by conjugating its elements are denoted, respectively,  $\mathbf{A}^T$ ,  $\mathbf{A}^*$ , and  $\bar{\mathbf{A}}$ . The spectral norm of a matrix  $\mathbf{A}$  will be denoted  $\|\mathbf{A}\|$ . If  $\mathbf{A}$  is square,  $\text{tr} \mathbf{A}$  refers to its trace. Let  $\mathbf{i} = \sqrt{-1}$ , then the operators  $\partial/\partial z$  and  $\partial/\partial \bar{z}$  where  $z = x + \mathbf{i}y$  is a complex number are defined by  $\frac{\partial}{\partial z} = \frac{1}{2} \left( \frac{\partial}{\partial x} - \mathbf{i} \frac{\partial}{\partial y} \right)$  and  $\frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial}{\partial x} + \mathbf{i} \frac{\partial}{\partial y} \right)$

where  $\frac{\partial}{\partial x}$  and  $\frac{\partial}{\partial y}$  are the standard partial derivatives with respect to  $x$  and  $y$ .

Throughout the paper, notation  $K$  will denote a generic constant whose main feature is *not* to depend on  $n$ . In particular, the value of  $K$  might change from a line to another as long as it never depends upon  $n$ . Constant  $K$  might depend on  $t \in \mathbb{R}^+$  and whenever needed, this dependence will be made more explicit. As usual, notation  $\alpha_n = \mathcal{O}(\beta_n)$  is a flexible shortcut for  $|\alpha_n| \leq K\beta_n$  and  $\alpha_n = o(\beta_n)$ , for  $\alpha_n = \varepsilon_n\beta_n$  with  $\varepsilon_n \rightarrow 0$  as  $n, N \rightarrow \infty$  (in the sense of (1)).

In order to study a deterministic approximation of  $I_n(\rho)$  and its fluctuations, the following mild assumptions are required over the two triangular arrays  $(d_i^{(n)}, 1 \leq i \leq N, n \geq 1)$  and  $(\tilde{d}_j^{(n)}, 1 \leq j \leq n, n \geq 1)$ .

**A1)** The real numbers  $d_i^{(n)}$  and  $\tilde{d}_j^{(n)}$  are nonnegative and the sequences  $(d_i^{(n)})$  and  $(\tilde{d}_j^{(n)})$  are uniformly bounded, i.e., there exist constants  $d_{\max}$  and  $\tilde{d}_{\max}$  such that

$$\sup_n \|\mathbf{D}_n\| < d_{\max} \text{ and } \sup_n \|\tilde{\mathbf{D}}_n\| < \tilde{d}_{\max}.$$

where  $\|\mathbf{D}_n\|$  and  $\|\tilde{\mathbf{D}}_n\|$  are the spectral norms of  $\mathbf{D}_n$  and  $\tilde{\mathbf{D}}_n$ .

**A2)** The normalized traces of  $\mathbf{D}_n$  and  $\tilde{\mathbf{D}}_n$  satisfy

$$\inf_n \frac{1}{n} \text{tr}(\mathbf{D}_n) > 0 \text{ and } \inf_n \frac{1}{n} \text{tr}(\tilde{\mathbf{D}}_n) > 0.$$

In the sequel, we shall frequently omit the subscript  $n$  and the superscript  $(n)$ .

The resolvent associated with  $\frac{1}{n}\mathbf{Y}_n\mathbf{Y}_n^*$  is the  $N \times N$  matrix  $\mathbf{H}_n(t) = (\frac{t}{n}\mathbf{Y}_n\mathbf{Y}_n^* + \mathbf{I}_N)^{-1}$ . Of prime importance is the random variable  $\beta(t) = \frac{1}{n}\text{tr}\mathbf{D}_n\mathbf{H}_n(t)$  and its expectation  $\alpha(t) = \frac{1}{n}\text{tr}\mathbf{D}_n\mathbb{E}\mathbf{H}_n(t)$ . We furthermore introduce the  $n \times n$  deterministic matrix defined by

$$\begin{aligned} \tilde{\mathbf{R}}_n(t) &= (\mathbf{I} + t\alpha(t)\tilde{\mathbf{D}}_n)^{-1} \\ &= \text{diag}(\tilde{r}_j(t), 1 \leq j \leq n) \\ \text{where } \tilde{r}_j(t) &= \frac{1}{1 + t\alpha(t)\tilde{d}_j} \end{aligned}$$

and the related quantity  $\tilde{\alpha}(t) = \frac{1}{n}\text{tr}\tilde{\mathbf{D}}_n\tilde{\mathbf{R}}_n(t)$ . In a symmetric fashion, the  $N \times N$  matrix  $\mathbf{R}_n(t)$  is defined by

$$\begin{aligned} \mathbf{R}_n(t) &= (\mathbf{I} + t\tilde{\alpha}(t)\mathbf{D}_n)^{-1} \\ &= \text{diag}(r_i(t), 1 \leq i \leq N) \text{ where } r_i(t) = \frac{1}{1 + t\tilde{\alpha}(t)d_i}. \end{aligned}$$

We finally introduce the solutions of a deterministic  $2 \times 2$  system.

*Proposition 1:* For every  $n$ , the system of equations in  $(\delta, \tilde{\delta})$

$$\begin{cases} \delta = \frac{1}{n}\text{tr}\mathbf{D}_n(\mathbf{I} + t\tilde{\delta}\mathbf{D}_n)^{-1} \\ \tilde{\delta} = \frac{1}{n}\text{tr}\tilde{\mathbf{D}}_n(\mathbf{I} + t\delta\tilde{\mathbf{D}}_n)^{-1} \end{cases} \quad (2)$$

admits a unique solution  $(\delta_n(t), \tilde{\delta}_n(t))$  satisfying  $\delta_n(t) > 0$ ,  $\tilde{\delta}_n(t) > 0$ . Moreover, there exist nonnegative measures  $\mu_n$  and  $\tilde{\mu}_n$  over  $\mathbb{R}^+$  such that

$$\delta_n(t) = \int_{\mathbb{R}^+} \frac{\mu_n(d\lambda)}{1 + t\lambda} \text{ and } \tilde{\delta}_n(t) = \int_{\mathbb{R}^+} \frac{\tilde{\mu}_n(d\lambda)}{1 + t\lambda} \quad (3)$$

where  $\mu_n(\mathbb{R}^+) = \frac{1}{n}\text{tr}\mathbf{D}_n$  and  $\tilde{\mu}_n(\mathbb{R}^+) = \frac{1}{n}\text{tr}\tilde{\mathbf{D}}_n$ .

Proof of Proposition 1 is postponed to the Appendix.

With  $\delta$  and  $\tilde{\delta}$  properly defined, we introduce the following  $N \times N$  and  $n \times n$  diagonal matrices:

$$\mathbf{T}_n = (\mathbf{I}_N + t\tilde{\delta}\mathbf{D}_n)^{-1} \text{ and } \tilde{\mathbf{T}}_n = (\mathbf{I}_n + t\delta\tilde{\mathbf{D}}_n)^{-1}.$$

Notice in particular that  $\delta = \frac{1}{n}\text{tr}\mathbf{D}_n\mathbf{T}_n$  and  $\tilde{\delta} = \frac{1}{n}\text{tr}\tilde{\mathbf{D}}_n\tilde{\mathbf{T}}_n$  by (2). We finally introduce the following quantities which are required to express the fluctuations of  $I_n(\rho)$ :

$$\begin{cases} \gamma_n(t) = \frac{1}{n}\text{tr}\mathbf{D}_n^2\mathbf{T}_n^2(t) \\ \tilde{\gamma}_n(t) = \frac{1}{n}\text{tr}\tilde{\mathbf{D}}_n^2\tilde{\mathbf{T}}_n^2(t). \end{cases} \quad (4)$$

*Proposition 2:* Assume that Assumptions **A1)** and **A2)** hold and denote by

$$\sigma_n^2(t) = -\log(1 - t^2\gamma_n(t)\tilde{\gamma}_n(t)), \quad t > 0 \quad (5)$$

where  $\gamma_n(t)$  and  $\tilde{\gamma}_n(t)$  are given by (4). Then  $\sigma_n^2(t)$  is well defined, i.e.,  $1 - t^2\gamma_n(t)\tilde{\gamma}_n(t) > 0$  for  $t > 0$ . Moreover, there exist nonnegative real numbers  $m_t$  and  $M_t$  such that

$$0 < m_t^2 \leq \inf_n \sigma_n^2(t) \leq \sup_n \sigma_n^2(t) \leq M_t^2 < \infty \text{ for } t > 0. \quad (6)$$

Finally,  $\sigma_n^2(t)$  is upper bounded uniformly in  $n$  and  $t$  for  $t \in [0, \rho]$ , i.e.,  $\sup_{t \leq \rho} M_t^2 < \infty$ .

Proof of Proposition 2 is postponed to the Appendix.

*1) Summary of the Main Notations:* In order to improve the readability of the paper, we gather all the notations in Table I. As expressed there, there are three kinds of quantities:

- 1) Random quantities,
- 2) Deterministic quantities depending on the law of  $\mathbf{Y}_n\mathbf{Y}_n^*$  via the expectation  $\mathbb{E}$  with respect to the entries of  $\mathbf{Y}_n$ ,
- 3) Deterministic quantities which only depend on the matrices  $\mathbf{D}_n$  and  $\tilde{\mathbf{D}}_n$ , sometimes via  $\delta$  and  $\tilde{\delta}$  (as defined in Proposition 1) which are easily computable.

The main goal of the forthcoming computations will be to approximate elements of the first and second kind by elements of the third kind.

### C. Statement of the Main Results

We now state the main results. Theorem 1 describes the first order approximation of the mutual information  $I_n(\rho)$  while Theorem 2 describes its fluctuations when centered with respect to its first order approximation.

TABLE I  
SUMMARY OF THE MAIN NOTATIONS

Random quantities	Deterministic quantities	
	depending on the law of $\mathbf{Y}\mathbf{Y}^*$ via $\mathbb{E}$	only depending on the variance structure via $\mathbf{D}$ and $\tilde{\mathbf{D}}$
$\mathbf{H} = \left(\frac{t}{n}\mathbf{Y}\mathbf{Y}^* + \mathbf{I}\right)^{-1}$ $\beta = \frac{1}{n}\text{tr}\mathbf{D}\mathbf{H}$	$\alpha = \frac{1}{n}\text{tr}\mathbf{D}(\mathbb{E}\mathbf{H})$ $\tilde{r}_j = (1 + t\alpha\tilde{d}_j)^{-1}$ $\tilde{\mathbf{R}} = (\mathbf{I} + t\alpha\tilde{\mathbf{D}})^{-1}$ $\tilde{\alpha} = \frac{1}{n}\text{tr}\tilde{\mathbf{D}}\tilde{\mathbf{R}} = \frac{1}{n}\text{tr}\tilde{\mathbf{D}}(\mathbf{I} + t\alpha\tilde{\mathbf{D}})^{-1}$ $r_i = (1 + t\tilde{\alpha}d_i)^{-1}$ $\mathbf{R} = (\mathbf{I} + t\tilde{\alpha}\mathbf{D})^{-1}$	$\delta = \frac{1}{n}\text{tr}\mathbf{D}(\mathbf{I} + t\tilde{\mathbf{D}})^{-1} = \frac{1}{n}\text{tr}\mathbf{D}\mathbf{T}$ $\tilde{\mathbf{T}} = (\mathbf{I} + t\tilde{\mathbf{D}})^{-1}$ $\tilde{\delta} = \frac{1}{n}\text{tr}\tilde{\mathbf{D}}(\mathbf{I} + t\tilde{\mathbf{D}})^{-1} = \frac{1}{n}\text{tr}\tilde{\mathbf{D}}\tilde{\mathbf{T}}$ $\mathbf{T} = (\mathbf{I} + t\tilde{\delta}\mathbf{D})^{-1}$ $\gamma = \frac{1}{n}\text{tr}\mathbf{T}^2\mathbf{D}^2, \quad \tilde{\gamma} = \frac{1}{n}\text{tr}\tilde{\mathbf{T}}^2\tilde{\mathbf{D}}^2$ $\sigma^2(t) = -\log(1 - t^2\gamma(t)\tilde{\gamma}(t))$

*Theorem 1:* Let  $\mathbf{X}_n$  be a  $N \times n$  matrix whose elements  $X_{ij}$  are independent complex Gaussian variables such that

$$\mathbb{E}(X_{ij}) = \mathbb{E}(X_{ij}^2) = 0, \quad \mathbb{E}(|X_{ij}|^2) = 1 \\ 1 \leq i \leq N, \quad 1 \leq j \leq n$$

and  $\mathbf{Y}_n = \mathbf{D}_n^{\frac{1}{2}}\mathbf{X}_n\tilde{\mathbf{D}}_n^{\frac{1}{2}}$  where the diagonal matrices  $\mathbf{D}_n$  and  $\tilde{\mathbf{D}}_n$  satisfy Assumptions **A1**) and **A2**). Let  $I_n(\rho) = \log \det \left( \frac{t}{n}\mathbf{Y}_n\mathbf{Y}_n^* + \mathbf{I}_N \right)$ . Then, we have

$$\mathbb{E}[I_n(\rho)] = V_n(\rho) + \mathcal{O}\left(\frac{1}{n}\right) \quad (7)$$

as  $n, N \rightarrow \infty$  (in the sense of (1)) where

$$V_n(\rho) = \log \det \left( \mathbf{I}_n + \rho\delta_n(\rho)\tilde{\mathbf{D}}_n \right) \\ + \log \det \left( \mathbf{I}_N + \rho\tilde{\delta}_n(\rho)\mathbf{D}_n \right) - n\rho\delta_n(\rho)\tilde{\delta}_n(\rho).$$

and where  $(\delta_n(t), \tilde{\delta}_n(t))$  is the unique positive solution of the system

$$\begin{cases} \delta = \frac{1}{n}\text{tr}\mathbf{D}_n(\mathbf{I}_N + t\tilde{\delta}\mathbf{D}_n)^{-1} \\ \tilde{\delta} = \frac{1}{n}\text{tr}\tilde{\mathbf{D}}_n(\mathbf{I}_n + t\delta\tilde{\mathbf{D}}_n)^{-1}. \end{cases}$$

*Theorem 2:* Assume that the setting of Theorem 1 holds and let

$$\sigma_n^2(\rho) = -\log(1 - \rho^2\gamma_n(\rho)\tilde{\gamma}_n(\rho))$$

where  $\gamma_n(\rho)$  and  $\tilde{\gamma}_n(\rho)$  are defined in (4). Then the following convergence holds true:

$$\frac{I_n(\rho) - V_n(\rho)}{\sigma_n(\rho)} \xrightarrow[n \rightarrow \infty]{\mathcal{L}} \mathcal{N}(0, 1)$$

where  $\xrightarrow{\mathcal{L}}$  stands for the convergence in distribution.

### III. MATHEMATICAL TOOLS AND SOME USEFUL RESULTS

In this section, we present the tools we will use extensively all along the paper. In Section III-A, we recall well-known matrix

results; in Section III-B, we present two fundamental properties of Gaussian models: The Integration by parts formula and Poincaré–Nash inequality for Gaussian vectors. Section III-C is devoted to a cornerstone approximation result which roughly states that  $\mathbf{R}$  and  $\tilde{\mathbf{R}}$  can be replaced with  $\mathbf{T}$  and  $\tilde{\mathbf{T}}$  up to some well-quantified error. In Section III-D, various variance estimates and approximation rules are stated.

#### A. General Results

1) *Some Matrix Inequalities:* Let  $\mathbf{A}$  and  $\mathbf{B}$  be two  $N \times N$  matrices with complex elements. Then

$$|\text{tr}(\mathbf{A}\mathbf{B})| \leq \sqrt{\text{tr}(\mathbf{A}\mathbf{A}^*)}\sqrt{\text{tr}(\mathbf{B}\mathbf{B}^*)}. \quad (8)$$

Assuming  $\mathbf{A}$  is Hermitian nonnegative, we have

$$|\text{tr}(\mathbf{A}\mathbf{B})| \leq \|\mathbf{B}\| \text{tr}(\mathbf{A}) \quad (9)$$

where  $\|\cdot\|$  is the spectral norm (see [26]).

2) *The Resolvent:* The Resolvent matrix  $\mathbf{H}_n(t)$  of matrix  $\mathbf{Y}_n\mathbf{Y}_n^*$  is defined as  $\mathbf{H}_n(t) = \left(\frac{t}{n}\mathbf{Y}_n\mathbf{Y}_n^* + \mathbf{I}_N\right)^{-1}$ . It is of constant use in this paper and we give here some of its properties. The following identity, also known as the *Resolvent identity*:

$$\mathbf{H}_n(t) = \mathbf{I}_N - \frac{t}{n}\mathbf{H}_n(t)\mathbf{Y}\mathbf{Y}^* \quad (10)$$

follows from the mere definition of  $\mathbf{H}_n$ . Furthermore, the spectral norm of the resolvent is readily bounded by one

$$\|\mathbf{H}_n(t)\| \leq 1 \text{ for } t \geq 0. \quad (11)$$

3) *Bounded Character of the Mean of Some Empirical Moments:* Let  $(\mathbf{B}_n)_{n \in \mathbb{N}} = \text{diag}\left(\left[b_1^{(n)}, \dots, b_n^{(n)}\right]\right)$ ,  $n \in \mathbb{N}$ , be a sequence of deterministic  $n \times n$  diagonal matrices. Assume **(A1)** and furthermore that  $\sup_n \|\mathbf{B}_n\| < \infty$ . Then, for every integer  $k$ , we have

$$\frac{1}{n}\mathbb{E}\left[\text{tr}\left(\frac{1}{n}\mathbf{Y}_n\mathbf{B}_n\mathbf{Y}_n^*\right)^k\right] < K_k. \quad (12)$$

Let us give a sketch of proof. Expanding the left-hand side of (12) yields

$$\frac{1}{n^{k+1}} \sum_{\substack{i_1, i_2, \dots, i_k=1:N \\ j_1, \dots, j_k=1:n}} b_{j_1} b_{j_2} \dots b_{j_k} \times \mathbb{E} \left[ Y_{i_1 j_1} \overline{Y_{i_2 j_2}} Y_{i_3 j_3} \overline{Y_{i_4 j_4}} \dots Y_{i_k j_k} \overline{Y_{i_1 j_k}} \right].$$

A close look at the argument of the  $\mathbb{E}$  operator implies that, due to the independence of the  $Y_{ij}$ , we only have  $k + 1$  degrees of freedom in the choice of the indices  $i_p$  and  $j_q$ . As all moments of the Gaussian law exist and as  $\|\mathbf{B}_n\|$ ,  $\|\mathbf{D}_n\|$ , and  $\|\tilde{\mathbf{D}}_n\|$  are bounded, this sum is of order 1 as  $n, N \rightarrow \infty$ .

4) *Differentiation Formulas*: Let  $\mathbf{A}$  be a  $N \times N$  complex matrix and let  $\mathbf{Q}(\mathbf{A}) = (\mathbf{I}_N + \mathbf{A})^{-1}$ . Let  $\delta\mathbf{A}$  be a perturbation of  $\mathbf{A}$ . Then

$$\mathbf{Q}(\mathbf{A} + \delta\mathbf{A}) = \mathbf{Q}(\mathbf{A}) - \mathbf{Q}(\mathbf{A})\delta\mathbf{A}\mathbf{Q}(\mathbf{A}) + o(\|\delta\mathbf{A}\|) \quad (13)$$

where  $o(\|\delta\mathbf{A}\|)$  is negligible with respect to  $\|\delta\mathbf{A}\|$  in a neighborhood of 0. Writing  $\mathbf{H}(t) = [H_{pq}(t)]_{p,q=1}^N$ , we need the expression of the partial derivative  $\partial H_{pq} / \partial Y_{ij}$ . Using (13), we have

$$\begin{aligned} \frac{\partial H_{pq}}{\partial Y_{ij}} &= -\frac{t}{n} \left[ \mathbf{H} \frac{\partial \mathbf{Y} \mathbf{Y}^*}{\partial Y_{ij}} \mathbf{H} \right]_{pq} \\ &= -\frac{t}{n} \left[ \mathbf{H} \left[ \delta(k-i) \overline{Y_{\ell j}} \right]_{k,\ell=1}^N \mathbf{H} \right]_{pq} \\ &= -\frac{t}{n} H_{pi} [\mathbf{Y}^* \mathbf{H}]_{jq} = -\frac{t}{n} H_{pi} [y_j^* \mathbf{H}]_q \end{aligned} \quad (14)$$

where  $\delta(\cdot)$  is the Kronecker function, i.e.,  $\delta(0) = 1$  otherwise  $\delta(\ell) = 0$ . Similarly, we can establish

$$\frac{\partial H_{pq}}{\partial \overline{Y_{ij}}} = -\frac{t}{n} [\mathbf{H} \mathbf{Y}]_{pj} H_{iq} = -\frac{t}{n} [\mathbf{H} y_j]_p H_{iq}. \quad (15)$$

The differential of  $g(\mathbf{A}) = \log \det(\mathbf{A})$  is given by  $g(\mathbf{A} + \delta\mathbf{A}) = g(\mathbf{A}) + \text{tr}(\mathbf{A}^{-1} \delta\mathbf{A}) + o(\|\delta\mathbf{A}\|)$ . We use this equation to derive the expression of  $\partial I(t) / \partial \overline{Y_{ij}}$  also needed below:

$$\begin{aligned} \frac{\partial I}{\partial \overline{Y_{ij}}} &= \frac{t}{n} \text{tr} \left( \mathbf{H} \frac{\partial \mathbf{Y} \mathbf{Y}^*}{\partial \overline{Y_{ij}}} \right) = \frac{t}{n} \text{tr} \left( \mathbf{H} \left[ \delta(\ell-j) Y_{k j} \right]_{k,\ell=1}^N \right) \\ &= \frac{t}{n} [\mathbf{H} \mathbf{Y}]_{ij} = \frac{t}{n} [\mathbf{H} y_j]_i. \end{aligned} \quad (16)$$

### B. Gaussian Tools

1) *An Integration by Parts Formula for Gaussian Functionals*: Let  $\boldsymbol{\xi} = [\xi_1, \dots, \xi_M]^T$  be a complex Gaussian random vector whose law is determined by  $\mathbb{E}[\boldsymbol{\xi}] = \mathbf{0}$ ,  $\mathbb{E}[\boldsymbol{\xi} \boldsymbol{\xi}^T] = \mathbf{0}$ , and  $\mathbb{E}[\boldsymbol{\xi} \boldsymbol{\xi}^*] = \boldsymbol{\Xi}$ . Let  $\Gamma = \Gamma(\xi_1, \dots, \xi_M, \overline{\xi_1}, \dots, \overline{\xi_M})$  be a  $C^1$  complex function polynomially bounded together with its derivatives, then

$$\mathbb{E} [\xi_p \Gamma(\boldsymbol{\xi})] = \sum_{m=1}^M [\boldsymbol{\Xi}]_{pm} \mathbb{E} \left[ \frac{\partial \Gamma(\boldsymbol{\xi})}{\partial \xi_m} \right]. \quad (17)$$

This formula relies on an integration by parts and thus is referred to as the Integration by parts formula for Gaussian vectors. It is

widely used in Mathematical Physics [27] and has been used in Random Matrix Theory in [17], [18].

2) *Poincaré–Nash Inequality*: Let  $\boldsymbol{\xi}$  and  $\Gamma$  be as previously and let  $\nabla_z \Gamma = [\partial \Gamma / \partial z_1, \dots, \partial \Gamma / \partial z_M]^T$  and  $\nabla_{\overline{z}} \Gamma = [\partial \Gamma / \partial \overline{z}_1, \dots, \partial \Gamma / \partial \overline{z}_M]^T$ . Then the following inequality holds true:

$$\begin{aligned} \text{var}(\Gamma(\boldsymbol{\xi})) &\leq \mathbb{E} \left[ \nabla_z \Gamma(\boldsymbol{\xi})^T \boldsymbol{\Xi} \nabla_{\overline{z}} \Gamma(\boldsymbol{\xi}) \right] \\ &\quad + \mathbb{E} \left[ (\nabla_{\overline{z}} \Gamma(\boldsymbol{\xi}))^* \boldsymbol{\Xi} \nabla_z \Gamma(\boldsymbol{\xi}) \right]. \end{aligned} \quad (18)$$

A proof of this inequality is available in [19] in the real case; see also [22]. When  $\boldsymbol{\xi}$  is the vector of the stacked columns of matrix  $\mathbf{Y}$ , i.e.,  $\boldsymbol{\xi} = [Y_{11}, \dots, Y_{Nn}]^T$ , formula (17) becomes

$$\mathbb{E} [Y_{ij} \Gamma(\mathbf{Y})] = d_i \tilde{d}_j \mathbb{E} \left[ \frac{\partial \Gamma(\mathbf{Y})}{\partial Y_{ij}} \right] \quad (19)$$

while inequality (18) writes

$$\text{var}(\Gamma(\mathbf{Y})) \leq \sum_{i=1}^N \sum_{j=1}^n d_i \tilde{d}_j \mathbb{E} \left[ \left| \frac{\partial \Gamma(\mathbf{Y})}{\partial Y_{i,j}} \right|^2 + \left| \frac{\partial \Gamma(\mathbf{Y})}{\partial \overline{Y_{i,j}}} \right|^2 \right]. \quad (20)$$

Poincaré–Nash inequality turns out to be extremely useful to deal with variances of various quantities of interest related to random matrices. In order to give right away the flavor of such results, we state and prove the following.

*Proposition 3*: Assume that the setting of Theorem 1 holds and let  $\mathbf{A}_n$  be a  $N \times N$  real diagonal matrix whose spectral norm is uniformly bounded in  $n$ . Then

$$\text{var} \left( \frac{1}{n} \text{tr} \mathbf{A}_n \mathbf{H}_n \right) = \mathcal{O}(n^{-2}).$$

*Proof*: We apply inequality (20) to the function  $\Gamma(\mathbf{Y}) = \frac{1}{n} \text{tr} \mathbf{A} \mathbf{H}$ . Using (14), we have

$$\frac{\partial \Gamma}{\partial Y_{i,j}} = \frac{1}{n} \sum_{p=1}^N a_p \frac{\partial H_{pp}}{\partial Y_{i,j}} = -\frac{t}{n^2} [y_j^* \mathbf{H} \mathbf{A} \mathbf{H}]_i.$$

Therefore, denoting by  $A$  the upper bound  $A = \sup_n \|\mathbf{A}_n\|$  and noticing that  $|\partial \Gamma / \partial Y_{i,j}| = |\partial \Gamma / \partial \overline{Y_{i,j}}|$ , we have:

$$\begin{aligned} \text{var} \Gamma(\mathbf{Y}) &\leq \frac{2t^2}{n^4} \sum_{i=1}^N \sum_{j=1}^n d_i \tilde{d}_j \mathbb{E} \left[ |y_j^* \mathbf{H} \mathbf{A} \mathbf{H}|_i^2 \right] \\ &= \frac{2t^2}{n^4} \sum_{j=1}^n \tilde{d}_j \mathbb{E} (y_j^* \mathbf{H} \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H} \mathbf{A} \mathbf{H} y_j) \\ &= \frac{2t^2}{n^3} \mathbb{E} \text{tr} \left( \mathbf{H} \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H} \mathbf{A} \mathbf{H} \frac{\mathbf{Y} \tilde{\mathbf{D}} \mathbf{Y}^*}{n} \right) \\ &\stackrel{(a)}{\leq} \frac{2t^2}{n^3} \mathbb{E} \left\{ \|\mathbf{H}\|^4 \|\mathbf{A}\|^2 \|\mathbf{D}\| \text{tr} \left( \frac{\mathbf{Y} \tilde{\mathbf{D}} \mathbf{Y}^*}{n} \right) \right\} \\ &\stackrel{(b)}{\leq} \frac{2A^2 d_{\max} t^2}{n^3} \mathbb{E} \text{tr} \left( \frac{\mathbf{Y} \tilde{\mathbf{D}} \mathbf{Y}^*}{n} \right) \stackrel{(c)}{\leq} \frac{K}{n^2}, \end{aligned}$$

where inequality (a) follows from (9), (b) follows from (11) and from the boundedness of  $\|\mathbf{A}_n\|$  and  $\|\mathbf{D}_n\|$ , and (c) follows from (12).  $\square$

### C. Approximation Rules

The following theorem is crucial in order to prove Theorems 1 and 2. Roughly speaking it allows to replace matrices  $\mathbf{R}_n$  and  $\tilde{\mathbf{R}}_n$  by  $\mathbf{T}_n$  and  $\tilde{\mathbf{T}}_n$  up to a well-quantified small error.

*Theorem 3:* Let  $(\mathbf{A}_n)$  and  $(\mathbf{B}_n)$  be two sequences of respectively  $N \times N$  and  $n \times n$  diagonal deterministic matrices whose spectral norm are uniformly bounded in  $n$ , then the following hold true:

$$\frac{1}{n} \text{tr} \mathbf{A}_n \mathbf{R}_n = \frac{1}{n} \text{tr} \mathbf{A}_n \mathbf{T}_n + \mathcal{O}\left(\frac{1}{n^2}\right) \quad (21)$$

$$\frac{1}{n} \text{tr} \mathbf{B}_n \tilde{\mathbf{R}}_n = \frac{1}{n} \text{tr} \mathbf{B}_n \tilde{\mathbf{T}}_n + \mathcal{O}\left(\frac{1}{n^2}\right). \quad (22)$$

Proof of Theorem 3 is postponed to the Appendix.

### D. More Variance Estimates and More Approximation Rules

We collect here a few results whose proofs rely on the Integration by parts formula (19), on Poincaré–Nash inequality, and on Theorem 3. The proofs of these results, although systematic, are somewhat lengthy and are therefore postponed to the Appendix. These results will be used extensively in Section V.

*Proposition 4:* In the setting of Theorem 1, let  $(\mathbf{A})$  and  $(\mathbf{B})$  be two sequences of respectively  $N \times N$  and  $n \times n$  diagonal deterministic matrices whose spectral norm are uniformly bounded in  $n$ . Consider the following functions:

$$\begin{aligned} \Phi(\mathbf{Y}) &= \frac{1}{n} \text{tr} \left( \mathbf{A} \mathbf{H} \frac{\mathbf{Y} \mathbf{B} \mathbf{Y}^*}{n} \right) \\ \Psi(\mathbf{Y}) &= \frac{1}{n} \text{tr} \left( \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H} \frac{\mathbf{Y} \mathbf{B} \mathbf{Y}^*}{n} \right). \end{aligned}$$

Then,

- 1) The following inequalities hold true:

$$\text{var}(\Phi(\mathbf{Y})) = \mathcal{O}(n^{-2}), \quad \text{var}(\Psi(\mathbf{Y})) = \mathcal{O}(n^{-2}).$$

- 2) The approximations hold true as shown in (23)–(24) at the bottom of the page.

#### IV. FIRST-ORDER MOMENT APPROXIMATION: PROOF OF THEOREM 1

This section is devoted to the proof of the following approximation:

$$\mathbb{E}[I_n(\rho)] = V_n(\rho) + \mathcal{O}(n^{-1}) \quad (25)$$

where

$$\begin{aligned} V_n(\rho) &= \log \det \left( \mathbf{I}_n + \rho \delta_n(\rho) \tilde{\mathbf{D}}_n \right) \\ &+ \log \det \left( \mathbf{I}_N + \rho \tilde{\delta}_n(\rho) \mathbf{D}_n \right) - n \rho \delta_n(\rho) \tilde{\delta}_n(\rho). \end{aligned} \quad (26)$$

This result already appears in [7] and is proved under greater generality in [9]. The proof presented here is new and relies on Gaussian tools.

*Outline of the Proof:* The proof is divided into three steps. We first make some preliminary remarks. Notice that the mutual information can be expressed as  $I(\rho) = \int_0^\rho \text{tr} (n^{-1} \mathbf{H}(t) \mathbf{Y} \mathbf{Y}^*) dt$ . In particular

$$\mathbb{E}[I(\rho)] = \int_0^\rho \text{tr} \left( \mathbb{E} \left[ \mathbf{H}(t) \frac{\mathbf{Y} \mathbf{Y}^*}{n} \right] \right) dt. \quad (27)$$

In order to study the asymptotic behavior of  $\mathbb{E}[I(\rho)]$ , it is thus enough to study  $\text{tr} \left( \mathbf{H}(t) \frac{\mathbf{Y} \mathbf{Y}^*}{n} \right)$  for  $n \rightarrow +\infty$  up to an integration. The Resolvent identity (10) yields

$$\text{tr} \mathbb{E} \left( \mathbf{H}(t) \frac{\mathbf{Y} \mathbf{Y}^*}{n} \right) = \text{tr} \mathbb{E} \left( \frac{\mathbf{I} - \mathbf{H}(t)}{t} \right).$$

We are therefore led to the study of  $\mathbb{E}[\text{tr}(\mathbf{H}(t))]$ . We now describe the three steps of the proof.

- A. In the first part of the proof, we expand  $\mathbb{E} \mathbf{H}(t)$  with the help of the Integration by parts formula (19). This derivation will bring to the fore the deterministic diagonal matrix  $\mathbf{R}$ , and Poincaré–Nash inequality will then allow us to obtain the following approximation:

$$\mathbb{E} \text{tr} \mathbf{A} \mathbf{H} = \text{tr} \mathbf{A} \mathbf{R} + \mathcal{O}(n^{-1})$$

for every diagonal matrix  $\mathbf{A}$  with a bounded spectral norm. Here are the main steps, gathered in an informal way. Differentiating the term  $\mathbb{E} \left( [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right)$ , we obtain

$$\begin{aligned} \mathbb{E} \left( [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right) &= d_p \tilde{d}_j \\ &\times \mathbb{E} [H_{pp}] - t \tilde{d}_j \mathbb{E} \left( \frac{1}{n} \text{tr}(\mathbf{D} \mathbf{H}) [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right) \end{aligned}$$

from which we will extract  $\mathbb{E}[H_{pp}]$  later on. At this point, Poincaré–Nash inequality yields some decorrelation up to  $\mathcal{O}(n^{-1})$  and we obtain

$$\begin{aligned} \mathbb{E} \left[ \frac{1}{n} \text{tr}(\mathbf{D} \mathbf{H}) (\mathbf{H} \mathbf{y}_j)_p \overline{Y_{pj}} \right] &\simeq \mathbb{E} \left[ \frac{1}{n} \text{tr}(\mathbf{D} \mathbf{H}) \right] \mathbb{E} \left[ [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right] \\ &= \alpha \mathbb{E} \left[ [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right]. \end{aligned}$$

This approximation allows us to isolate  $\mathbb{E} \left( [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right)$  shown in the equation at the bottom of the following page.

Now summing over  $j$  and using the Resolvent identity  $\mathbb{E} H_{pp} = 1 - \frac{t}{n} \sum_{j=1}^n \mathbb{E} [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}}$  in the previous equation yields

$$\frac{1 - \mathbb{E} H_{pp}}{t} \simeq \tilde{\alpha} d_p \mathbb{E} H_{pp}, \quad \text{that is } \mathbb{E} H_{pp} \simeq r_p.$$

All the technical details are provided in Section IV-A.

$$\mathbb{E}[\Phi(\mathbf{Y})] = \frac{1}{n} \text{tr} \left( \tilde{\mathbf{D}} \tilde{\mathbf{T}} \mathbf{B} \right) \frac{1}{n} \text{tr}(\mathbf{A} \mathbf{D} \mathbf{T}) + \mathcal{O}(n^{-2}) \quad (23)$$

$$\mathbb{E}[\Psi(\mathbf{Y})] = \frac{1}{1 - t^2 \gamma \tilde{\gamma}} \left( \frac{1}{n^2} \text{tr} \left( \tilde{\mathbf{D}} \tilde{\mathbf{T}} \mathbf{B} \right) \text{tr}(\mathbf{A} \mathbf{D}^2 \mathbf{T}^2) - \frac{t \gamma}{n^2} \text{tr} \left( \tilde{\mathbf{D}}^2 \tilde{\mathbf{T}}^2 \mathbf{B} \right) \text{tr}(\mathbf{A} \mathbf{D} \mathbf{T}) \right) + \mathcal{O}\left(\frac{1}{n^2}\right). \quad (24)$$

B. The second step follows from the approximation rule (21) stated in Section III-C, which immediately yields

$$\mathbb{E} \text{tr} \mathbf{A} \mathbf{H} = \text{tr} \mathbf{A} \mathbf{T} + \mathcal{O}(n^{-1}).$$

This in turn will imply that

$$\begin{aligned} \mathbb{E} \text{tr} \left( \mathbf{H}(t) \frac{\mathbf{Y} \mathbf{Y}^*}{n} \right) &= \text{tr} \left( \frac{\mathbf{I} - \mathbb{E} \mathbf{H}}{t} \right) = \text{tr} \left( \frac{\mathbf{I} - \mathbf{T}}{t} \right) + \varepsilon_n(t) \\ &\stackrel{(a)}{=} n \delta(t) \tilde{\delta}(t) + \varepsilon_n(t). \end{aligned}$$

where (a) follows from the fact that  $\mathbf{I} - \mathbf{T} = t \tilde{\delta} \mathbf{D} (\mathbf{I} + t \tilde{\delta} \mathbf{D})^{-1}$ .

C. In the third step, we integrate the previous equality:

$$\int_0^\rho \mathbb{E} \text{tr} \left( \mathbf{H}(t) \frac{\mathbf{Y} \mathbf{Y}^*}{n} \right) dt = n \int_0^\rho \delta(t) \tilde{\delta}(t) dt + \int_0^\rho \varepsilon_n(t) dt.$$

We identify  $n \int_0^\rho \delta(t) \tilde{\delta}(t) dt$  with  $V_n(\rho)$  as given by (26), and check that  $\int_0^\rho \varepsilon_n(t) dt = \mathcal{O}(n^{-1})$ .

A. *Development of  $\mathbb{E}(\text{tr} \mathbf{A} \mathbf{H}(t))$  and Approximation by  $\text{tr} \mathbf{A} \mathbf{R}(t)$*

In order to study  $\mathbb{E}(\text{tr} \mathbf{A} \mathbf{H}(t))$ , we first consider the diagonal entries  $H_{pp}(t)$  of  $\mathbf{H}(t)$ . For each index  $j$ , we have

$$\mathbb{E} \left( [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right) = \sum_{i=1}^N \mathbb{E} (H_{pi} Y_{ij} \overline{Y_{pj}}).$$

We now apply the Integration by parts formula (19) to the summand of the right-hand side for function  $\Gamma$  defined as  $\Gamma(\mathbf{Y}) = H_{pi} \overline{Y_{pj}}$ . This yields

$$\begin{aligned} \mathbb{E} (H_{pi} Y_{ij} \overline{Y_{pj}}) &= d_i \tilde{d}_j \mathbb{E} [H_{ii}] \delta(i - p) \\ &\quad - d_i \tilde{d}_j \frac{t}{n} \mathbb{E} \left( [\mathbf{H} \mathbf{y}_j]_p H_{ii} \overline{Y_{pj}} \right). \end{aligned} \quad (28)$$

Therefore

$$\begin{aligned} \mathbb{E} \left( [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right) &= d_p \tilde{d}_j \mathbb{E} [H_{pp}] - t \tilde{d}_j \\ &\quad \times \mathbb{E} \left( \frac{1}{n} \text{tr}(\mathbf{D} \mathbf{H}) [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right) \end{aligned} \quad (29)$$

from which we extract  $\mathbb{E}[H_{pp}]$  later on. Recall at this point that  $\text{var}(n^{-1} \text{tr} \mathbf{D} \mathbf{H}(t)) = \mathcal{O}(n^{-2})$  by Proposition 3. Recall also the following notations:  $\beta = n^{-1} \text{tr}(\mathbf{D} \mathbf{H})$ ,  $\alpha = \mathbb{E}[\beta]$ , and  $\hat{\beta} = \beta - \alpha$ . Plugging the relation  $\beta = \alpha + \hat{\beta}$  into (29), we get

$$\begin{aligned} \mathbb{E} \left[ [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right] &= d_p \tilde{d}_j \mathbb{E} [H_{pp}] \\ &\quad - t \tilde{d}_j \alpha \mathbb{E} \left[ [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right] - t \tilde{d}_j \mathbb{E} \left[ \hat{\beta} [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right]. \end{aligned} \quad (30)$$

Solving this equation w.r.t.  $\mathbb{E} \left[ [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right]$  provides

$$\mathbb{E} \left[ [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right] = d_p \tilde{d}_j \tilde{r}_j \mathbb{E} [H_{pp}] - t \tilde{d}_j \tilde{r}_j \mathbb{E} \left[ \hat{\beta} [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right]$$

$$\text{where } \tilde{r}_j(t) = \frac{1}{1 + t \alpha(t) \tilde{d}_j} \text{ for } 1 \leq j \leq n. \quad (31)$$

Summing (31) over  $j$  yields

$$\mathbb{E} \left[ \mathbf{H} \frac{\mathbf{Y} \mathbf{Y}^*}{n} \right]_{pp} = \tilde{\alpha} d_p \mathbb{E} [H_{pp}] - t \mathbb{E} \hat{\beta} \left[ \mathbf{H} \frac{\mathbf{Y} \tilde{\mathbf{D}} \tilde{\mathbf{R}} \mathbf{Y}^*}{n} \right]_{pp} \quad (32)$$

where  $\tilde{\mathbf{R}}$  is the diagonal matrix  $\text{diag}(\tilde{r}_j(t)) = (\mathbf{I} + \alpha t \tilde{\mathbf{D}})^{-1}$  and  $\tilde{\alpha} = \frac{1}{n} \text{tr} \tilde{\mathbf{D}} \tilde{\mathbf{R}}$ . In order to obtain an expression for  $\mathbb{E}[H_{pp}]$ , we plug the identity (32) into the Resolvent identity

$$\mathbb{E} [H_{pp}] = 1 - t \mathbb{E} \left[ \left[ \mathbf{H} \frac{\mathbf{Y} \mathbf{Y}^*}{n} \right]_{pp} \right]$$

and obtain

$$\mathbb{E} [H_{pp}] = r_p + t^2 r_p \mathbb{E} \left[ \hat{\beta} \left[ \mathbf{H} \frac{\mathbf{Y} \tilde{\mathbf{D}} \tilde{\mathbf{R}} \mathbf{Y}^*}{n} \right]_{pp} \right] \quad (33)$$

with  $r_p(t) = (1 + t \tilde{\alpha} d_p)^{-1}$ . Let  $\mathbf{A}$  be a  $N \times N$  diagonal matrix with bounded spectral norm. Multiplying (33) by  $\mathbf{A}$ 's components and summing over  $p$  yields

$$\mathbb{E} \text{tr}(\mathbf{A} \mathbf{H}) = \text{tr}(\mathbf{A} \mathbf{R}) + n t^2 \mathbb{E} \left[ \hat{\beta} \Phi(\mathbf{Y}) \right]$$

where  $\Phi(\mathbf{Y}) = \frac{1}{n} \text{tr}(\mathbf{A} \mathbf{R} \mathbf{H} \frac{\mathbf{Y} \tilde{\mathbf{D}} \tilde{\mathbf{R}} \mathbf{Y}^*}{n})$ . As  $\hat{\beta}$  is zero-mean,  $\mathbb{E}[\hat{\beta} \Phi] = \mathbb{E}[\hat{\beta} \hat{\Phi}]$ . In particular, Cauchy-Schwarz inequality yields

$$|\mathbb{E}[\hat{\beta} \hat{\Phi}]| \leq \sqrt{\text{var}(\hat{\beta})} \sqrt{\text{var}(\Phi)}.$$

Recall that  $\text{var}(\beta) = \mathcal{O}(n^{-2})$  by Proposition 3. Since  $\|\mathbf{R}\|$  and  $\|\tilde{\mathbf{D}} \tilde{\mathbf{R}}\|$  are both bounded by Assumption **A1** and by the definitions of  $\mathbf{R}$  and  $\tilde{\mathbf{R}}$ , one can directly apply the result of Proposition 4 to  $\Phi$  in order to get  $\text{var}(\Phi) = \mathcal{O}(n^{-2})$ .

We have, therefore, proved the following proposition.

*Proposition 5:* In the setting of Theorem 1, let  $\mathbf{A}_n$  be a uniformly bounded diagonal  $N \times N$  matrix. Then for every  $t \in \mathbb{R}^+$ ,

$$\mathbb{E}(\text{tr} \mathbf{A}_n \mathbf{H}_n(t)) = \text{tr} \mathbf{A}_n \mathbf{R}_n(t) + \mathcal{O}(n^{-1}). \quad (34)$$

B. *The Deterministic Approximation  $\mathbf{T}_n(t)$*

Proposition 5 provides a deterministic equivalent to  $\mathbb{E}(\text{tr} \mathbf{A}_n \mathbf{H}_n)$  since the matrix  $\mathbf{R}_n$  is deterministic; however its elements still depend on  $\tilde{\alpha} = n^{-1} \text{tr}(\tilde{\mathbf{D}}_n \tilde{\mathbf{R}}_n)$ , which itself depends on  $\alpha = \mathbb{E}(n^{-1} \text{tr} \mathbf{D}_n \mathbf{H}_n)$ , an unknown parameter. The next step is therefore to apply Theorem 3 to approximate the matrix  $\mathbf{R}_n$  by  $\mathbf{T}_n$ , which only depends on  $\mathbf{D}_n$  and  $\tilde{\mathbf{D}}_n$  and

$$(1 + t \tilde{d}_j \alpha) \mathbb{E} \left( [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right) \simeq d_p \tilde{d}_j \mathbb{E} [H_{pp}] \Leftrightarrow \mathbb{E} \left( [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} \right) \simeq d_p \tilde{d}_j \tilde{r}_j \mathbb{E} [H_{pp}].$$

on  $\delta$  and  $\tilde{\delta}$ , the solutions of (2). Theorem 3 together with (34) imply that:

$$\mathbb{E}(\text{tr} \mathbf{A}_n \mathbf{H}_n) = \text{tr}(\mathbf{A}_n \mathbf{T}_n) + \mathcal{O}(n^{-1}). \quad (35)$$

Since  $\mathbf{T}_n$  only depends on  $\delta$  and  $\tilde{\delta}$ , (35) provides a deterministic equivalent of  $\mathbb{E}(\text{tr} \mathbf{A}_n \mathbf{H}_n)$  in terms of  $\delta$  and  $\tilde{\delta}$ . Note that taking  $\mathbf{A} = \mathbf{D}$  yields in particular  $\alpha = \delta + \mathcal{O}(n^{-2})$  while a direct application of Theorem 3 for  $\tilde{\mathbf{A}}_n = \tilde{\mathbf{D}}_n$  yields  $\tilde{\alpha} = \tilde{\delta} + \mathcal{O}(n^{-2})$ .

We are now in a position to describe the behavior of  $\mathbb{E} \text{tr} \left( \mathbf{H}_n(t) \frac{\mathbf{Y}_n \mathbf{Y}_n^*}{n} \right)$  by using the Resolvent identity. From (10) and (35), taking  $\mathbf{A}_n = \mathbf{I}_n$ , we immediately obtain

$$\mathbb{E} \text{tr} \left( \mathbf{H}_n(t) \frac{\mathbf{Y}_n \mathbf{Y}_n^*}{n} \right) = \frac{1}{t} \text{tr}(\mathbf{I}_n - \mathbf{T}_n(t)) + \mathcal{O}(n^{-1}).$$

As  $\mathbf{I}_n - \mathbf{T}_n(t) = (\mathbf{T}_n(t)^{-1} - \mathbf{I}_n) \mathbf{T}_n(t) = t \tilde{\delta}(t) \mathbf{D}_n \mathbf{T}_n(t)$ , we eventually get that

$$\mathbb{E} \left[ \text{tr} \left( \mathbf{H}_n(t) \frac{\mathbf{Y}_n \mathbf{Y}_n^*}{n} \right) \right] = n \delta(t) \tilde{\delta}(t) + \varepsilon_n(t) \quad (36)$$

where the error  $\varepsilon_n(t)$  is a  $\mathcal{O}(n^{-1})$  term.

### C. Recovering the Deterministic Approximation $V(\rho)$

As mentioned previously,  $\varepsilon_n(t)$  is a  $\mathcal{O}(n^{-1})$  term, i.e.,  $|\varepsilon_n(t)| \leq K_t n^{-1}$ . One can easily keep track of  $K_t$  in the derivations that lead to (36) and prove that  $K_t$  is bounded on the compact interval  $[0, \rho]$ . In particular,  $|\varepsilon_n(t)| < K n^{-1}$  on the compact interval  $[0, \rho]$  for some  $K > 0$ . The proof of this fact is omitted.

As  $n \varepsilon_n(t)$  is uniformly bounded on  $[0, \rho]$ , we have  $\left| \int_0^\rho \varepsilon_n(t) dt \right| = \mathcal{O}(n^{-1})$ . Therefore

$$\mathbb{E}[I(\rho)] = \int_0^\rho n \delta(t) \tilde{\delta}(t) dt + \mathcal{O}(n^{-1}).$$

Consider now

$$V(\rho) = W(\rho, \delta(\rho), \tilde{\delta}(\rho))$$

where function  $W(\rho, \delta, \tilde{\delta})$  is defined by

$$W(\rho, \delta, \tilde{\delta}) = \log \det(\mathbf{I}_n + \rho \delta \tilde{\mathbf{D}}_n) + \log \det(\mathbf{I}_n + \rho \tilde{\delta} \mathbf{D}_n) - n \rho \delta \tilde{\delta}.$$

One can easily check that

$$\begin{aligned} \frac{\partial W}{\partial \delta} &= \rho \left( \text{tr} \left( \tilde{\mathbf{D}}_n (\mathbf{I}_n + \rho \delta \tilde{\mathbf{D}}_n)^{-1} \right) - n \tilde{\delta} \right) \text{ and} \\ \frac{\partial W}{\partial \tilde{\delta}} &= \rho \left( \text{tr} \left( \mathbf{D}_n (\mathbf{I}_n + \rho \tilde{\delta} \mathbf{D}_n)^{-1} \right) - n \delta \right). \end{aligned}$$

As the pair  $(\delta(\rho), \tilde{\delta}(\rho))$  satisfies (2), the above partial derivatives evaluated at point  $(\rho, \delta(\rho), \tilde{\delta}(\rho))$  are zero. Therefore

$$\frac{dV}{d\rho} = \left( \frac{\partial W}{\partial \rho} \right)_{(\rho, \delta(\rho), \tilde{\delta}(\rho))} = n \delta(\rho) \tilde{\delta}(\rho) \quad (37)$$

which in turn implies (7). Theorem 1 is proved.

*Remark 1 (on the Deterministic Approximation  $\mathbf{T}_n$ ):* The deterministic approximation  $\mathbf{T}_n$  can be used to approximate func-

tions of the eigenvalues of  $\mathbf{Y}_n \mathbf{Y}_n^*$  other than the mutual information  $\log \det(\rho n^{-1} \mathbf{Y}_n \mathbf{Y}_n^* + \mathbf{I}_n)$  (see for instance [9]). This relies on a specific representation of  $\mathbf{T}_n$ . The spectral theorem for Hermitian matrices yields the integral representation:

$$\frac{1}{n} \text{tr} \mathbf{H}_n(z) = \int_0^\infty \frac{N_n(d\lambda)}{1 + \lambda z}, \quad z \in \mathbb{C} \setminus \mathbb{R}_-$$

where  $N_n$  represents the empirical distribution of the eigenvalues of  $\mathbf{Y}_n \mathbf{Y}_n^*$ . It can be shown that  $n^{-1} \text{tr} \mathbf{T}_n$  admits a similar representation:

$$\frac{1}{n} \text{tr} \mathbf{T}_n(z) = \int_0^\infty \frac{\pi_n(d\lambda)}{1 + \lambda z}, \quad z \in \mathbb{C} \setminus \mathbb{R}_-$$

where  $\pi_n$  is a probability measure. Finally, one can prove that  $\int_0^\infty f(\lambda) N_n(d\lambda) - \int_0^\infty f(\lambda) \pi_n(d\lambda)$  converges to zero almost surely for every continuous bounded function (see [9], for details).

## V. SECOND-ORDER ANALYSIS: PROOF OF THEOREM 2

This section is devoted to the proof of the Central Limit Theorem

$$\sigma_n^{-1}(\rho) (I_n(\rho) - V_n(\rho)) \xrightarrow[n, N \rightarrow \infty]{\mathcal{L}} \mathcal{N}(0, 1). \quad (38)$$

Denote by  $\psi_n(u, \rho) = \mathbb{E} [e^{iu(I_n(\rho) - V_n(\rho))}]$  the characteristic function of  $I_n(\rho) - V_n(\rho)$ . We first reduce the problem in the following way.

*Proposition 6:* Assume that for every  $u \in \mathbb{R}$ ,

$$h_n(u) = \psi_n(u, \rho) - e^{-u^2 \sigma_n^2(\rho)/2} \xrightarrow[n, N \rightarrow \infty]{} 0 \quad (39)$$

then (38) holds true.

*Proof:*<sup>2</sup> We first prove that the sequence  $\sigma_n^{-1} (I_n - V_n)$  is tight (we drop the dependence in  $\rho$ ). Recall from Proposition 2 that  $0 < m \leq \sigma_n^2 \leq M < \infty$ . Let  $\varepsilon > 0$ . For  $x > 0$  small enough, we have

$$\frac{1}{x} \int_{-x}^x (1 - e^{-u^2 \sigma_n^2/2}) du \leq \frac{1}{x} \int_{-x}^x (1 - e^{-u^2 M/2}) du \leq \varepsilon.$$

Moreover, since  $\psi_n(u) - e^{-u^2 \sigma_n^2} \rightarrow 0$ , the Dominated Convergence Theorem yields

$$\left| \frac{1}{x} \int_{-x}^x (1 - \psi_n(u)) du - \frac{1}{x} \int_{-x}^x (1 - e^{-u^2 \sigma_n^2}) du \right| \leq \varepsilon$$

for  $n$  large enough. Now, given a real random variable  $X$  with characteristic function  $\varphi(u)$ , the following inequality holds true:

$$\mathbb{P} \left[ |X| \geq \frac{2}{x} \right] \leq \frac{1}{x} \int_{-x}^x (1 - \varphi(u)) du$$

(see, for instance, [28, eq. (26.22), Theorem 26.3]). Applying this inequality to  $I_n - V_n$ , we obtain

$$\mathbb{P} [|I_n - V_n| \geq 2/x] \leq 2\varepsilon$$

<sup>2</sup>This proof simplifies an earlier proof by the authors and was suggested by one of the reviewers whom the authors would like to thank.



for  $n$  large enough; in other words,  $I_n - V_n$  is tight. Tightness of  $\sigma_n^{-1} (I_n - V_n)$  follows from the fact that  $\sigma_n^2 \geq m > 0$ . We are now in position to conclude. Since  $\sigma_n^{-1} (I_n - V_n)$  is tight, one can extract a subsequence  $(n)$  such that  $\sigma_n^{-1} (I_n - V_n)$  converges. Now, since  $\sigma_n$  belongs to a compact set, one can extract a converging subsequence from  $(n)$ , say  $(m)$ , such that  $\sigma_m^2 \rightarrow a > 0$ . From (39), we have  $I(m) - V(m) \xrightarrow{\mathcal{L}} \mathcal{N}(0, a)$  or equivalently  $\sigma_m^{-1} (I(m) - V(m)) \xrightarrow{\mathcal{L}} \mathcal{N}(0, 1)$ , and the limit of  $\sigma_n^{-1} (I_n - V_n)$  (which is the same as the one of the subsequence  $\sigma_m^{-1} (I(m) - V(m))$ ) is necessarily  $\mathcal{N}(0, 1)$ .

We have proved that for every subsequence  $(n)$  such that  $\sigma_n^{-1}(\rho) (I_n(\rho) - V_n(\rho))$  converges,

$$\sigma_n^{-1}(\rho) (I_n(\rho) - V_n(\rho)) \xrightarrow{\mathcal{L}} \mathcal{N}(0, 1).$$

The result remains true for the whole sequence  $\sigma_n^{-1} (I_n - V_n)$  by Corollary of [28, Theorem 25.10]  $\square$

*Outline of the Proof of (39):* The proof of the convergence of  $h_n(u)$  toward zero is divided into two steps.

A. We first differentiate  $\psi_n(u, t)$  with respect to  $t$  in order to obtain a differential equation of the form:

$$\frac{\partial \psi_n(u, t)}{\partial t} = -\frac{u^2}{2} \eta_n(t) \psi_n(u, t) + \varepsilon_n(u, t). \quad (40)$$

In order to obtain the differential (40), we first develop  $\partial \psi / \partial t$  with the help of the Integration by parts formula (19). We then use Poincaré–Nash inequality to prove that relevant variances are of order  $\mathcal{O}(n^{-2})$ . This will enable us to decorrelate various expectations, i.e., to express them as products of expectations up to negligible terms. We shall then use the approximation rules stated in Proposition 4 in Section III-D to deal with the obtained expectations.

B. The second step is devoted to identify the variance, that is to prove the identity

$$\int_0^\rho \eta_n(t) dt = \sigma_n^2(\rho)$$

where  $\sigma_n^2$  is given by (5), i.e.,  $\sigma_n^2(\rho) = -\log(1 - \rho^2 \gamma_n(\rho) \tilde{\gamma}_n(\rho))$ .

C. The third step is devoted to the integration of (40). Instead of directly integrating (40), we introduce  $K_n(u, \rho) = \psi_n(u, \rho) e^{\frac{u^2}{2} \sigma_n^2(\rho)}$  which satisfies the following differential equation:

$$\frac{\partial K_n(u, t)}{\partial t} = \varepsilon_n(u, t) e^{\frac{u^2}{2} \sigma_n^2(t)}. \quad (41)$$

Taking into account the obvious facts that  $\psi_n(u, 0) = 1$ ,  $\sigma_n^2(0) = 0$  and therefore that  $K_n(u, 0) = 1$ , we shall obtain that

$$K_n(u, \rho) = 1 + \int_0^\rho \varepsilon_n(u, t) e^{\frac{u^2}{2} \sigma_n^2(t)} dt$$

and prove that  $\int_0^\rho \varepsilon_n(u, t) e^{\frac{u^2}{2} \sigma_n^2(t)} dt = \mathcal{O}(n^{-1})$ . This will yield in turn that

$$\begin{aligned} \psi_n(u, \rho) &= (1 + \mathcal{O}(n^{-1})) e^{-\frac{u^2}{2} \sigma_n^2(\rho)} \\ &\stackrel{(a)}{=} e^{-\frac{u^2}{2} \sigma_n^2(\rho)} + \mathcal{O}(n^{-1}), \end{aligned}$$

where (a) follows from Proposition 2.

The theorem will then be proved.

A. *The Differential Equation*  $\partial_t \psi_n = -\frac{u^2}{2} \eta_n \psi_n + \varepsilon_n$

Recall that  $\psi_n(u, \rho) = \varphi_n(u, \rho) e^{-\mathbf{i}uV_n(\rho)}$  where  $\varphi(u, t) = \mathbb{E} (e^{\mathbf{i}uI(t)})$ . As  $V_n'(t) = n\delta(t)\delta(t)$  by (37), we obtain

$$\frac{\partial \psi(u, t)}{\partial t} = e^{-\mathbf{i}uV(t)} \frac{\partial \varphi(u, t)}{\partial t} - \mathbf{i}un\delta(t)\delta(t)\psi(u, t). \quad (42)$$

Since  $I'(t) = n^{-1} \text{tr} \mathbf{H}(t) \mathbf{Y} \mathbf{Y}^*$  by (27), we have:

$$\begin{aligned} \frac{\partial \varphi(u, t)}{\partial t} &= \mathbf{i}u \mathbb{E} \left[ \text{tr} \left( \mathbf{H}(t) \frac{\mathbf{Y} \mathbf{Y}^*}{n} \right) e^{\mathbf{i}uI(t)} \right] \\ &= \frac{\mathbf{i}u}{n} \sum_{p,i=1}^N \sum_{j=1}^n \mathbb{E} [Y_{ij} H_{pi} \overline{Y_{pj}} e^{\mathbf{i}uI}]. \end{aligned} \quad (43)$$

Applying the Integration by parts formula (19) to  $\mathbb{E} [Y_{ij} H_{pi} \overline{Y_{pj}} e^{\mathbf{i}uI}]$  (which can be written  $\mathbb{E} (Y_{ij} \Gamma(\mathbf{Y}))$  for  $\Gamma(\mathbf{Y}) = H_{pi} \overline{Y_{pj}} e^{\mathbf{i}uI}$ ) and using the differentiation formulas (15) and (16) yields

$$\begin{aligned} \mathbb{E} [Y_{ij} H_{pi} \overline{Y_{pj}} e^{\mathbf{i}uI}] &= d_i \tilde{d}_j \mathbb{E} \left[ \frac{\partial}{\partial Y_{ij}} (H_{pi} \overline{Y_{pj}} e^{\mathbf{i}uI}) \right] \\ &= -\frac{t}{n} d_i \tilde{d}_j \mathbb{E} \left[ [\mathbf{H} \mathbf{y}_j]_p H_{ii} \overline{Y_{pj}} e^{\mathbf{i}uI} \right] \\ &\quad + d_i \tilde{d}_j \delta(i-p) \mathbb{E} [H_{pi} e^{\mathbf{i}uI}] \\ &\quad + \frac{\mathbf{i}ut}{n} d_i \tilde{d}_j \mathbb{E} [H_{pi} \overline{Y_{pj}} [\mathbf{H} \mathbf{y}_j]_i e^{\mathbf{i}uI}]. \end{aligned} \quad (44)$$

We now sum over index  $i$  and obtain

$$\begin{aligned} \mathbb{E} \left[ [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} e^{\mathbf{i}uI} \right] &= -t \tilde{d}_j \mathbb{E} \left[ \beta [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} e^{\mathbf{i}uI} \right] \\ &\quad + d_p \tilde{d}_j \mathbb{E} [H_{pp} e^{\mathbf{i}uI}] \\ &\quad + \frac{\mathbf{i}ut}{n} \tilde{d}_j \mathbb{E} \left[ [\mathbf{H} \mathbf{D} \mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} e^{\mathbf{i}uI} \right] \end{aligned}$$

where  $\beta = n^{-1} \text{tr} \mathbf{D} \mathbf{H}$ . Writing  $\beta = \hat{\beta} + \alpha$  yields

$$\begin{aligned} (1 + t\alpha \tilde{d}_j) \mathbb{E} \left[ [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} e^{\mathbf{i}uI} \right] &= -t \tilde{d}_j \mathbb{E} \left[ \hat{\beta} [\mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} e^{\mathbf{i}uI} \right] \\ &\quad + d_p \tilde{d}_j \mathbb{E} [H_{pp} e^{\mathbf{i}uI}] + \frac{\mathbf{i}ut}{n} \tilde{d}_j \mathbb{E} \left[ [\mathbf{H} \mathbf{D} \mathbf{H} \mathbf{y}_j]_p \overline{Y_{pj}} e^{\mathbf{i}uI} \right]. \end{aligned} \quad (45)$$

We now take into account that  $\tilde{r}_j(t) = (1 + t\alpha \tilde{d}_j)^{-1}$  and sum over  $j$ : see (46) at the top of the following page.

By the Resolvent identity (10),  $\mathbb{E} [H_{pp} e^{\mathbf{i}uI}] = \mathbb{E} [e^{\mathbf{i}uI}] - \frac{t}{n} \mathbb{E} \left[ [\mathbf{H} \mathbf{Y} \mathbf{Y}^*]_{pp} e^{\mathbf{i}uI} \right]$ . Replace now in (46), recall that  $r_p(t) = (1 + t\tilde{\alpha}(t) d_p)^{-1}$  and sum over  $p$  to obtain

$$\begin{aligned} &\mathbb{E} \left[ \text{tr} \left( \mathbf{H} \frac{\mathbf{Y} \mathbf{Y}^*}{n} \right) e^{\mathbf{i}uI} \right] \\ &= \text{tr} (\mathbf{D} \mathbf{R}) \hat{\alpha} \mathbb{E} [e^{\mathbf{i}uI}] \end{aligned}$$

$$\begin{aligned} \mathbb{E} \left[ [\mathbf{H}\mathbf{Y}\mathbf{Y}^*]_{pp} e^{iuI} \right] &= -t \mathbb{E} \left[ \hat{\beta} \left[ \mathbf{H}\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\mathbf{Y}^* \right]_{pp} e^{iuI} \right] \\ &\quad + n\hat{\alpha}d_p \mathbb{E} [H_{pp}e^{iuI}] + \frac{iut}{n} \mathbb{E} \left[ [\mathbf{H}\mathbf{D}\mathbf{H}\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\mathbf{Y}^*]_{pp} e^{iuI} \right]. \end{aligned} \quad (46)$$

$$\begin{aligned} &+ iut \mathbb{E} \left[ \frac{1}{n} \text{tr} \left( \mathbf{R}\mathbf{H}\mathbf{D}\mathbf{H} \frac{\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\mathbf{Y}^*}{n} \right) e^{iuI} \right] \\ &= -t \mathbb{E} \left[ \hat{\beta} \text{tr} \left( \mathbf{R}\mathbf{H} \frac{\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\mathbf{Y}^*}{n} \right) e^{iuI} \right] \\ &\triangleq \chi_1 + \chi_2 + \chi_3. \end{aligned} \quad (47)$$

Thanks to Theorem 3

$$\begin{aligned} \chi_1 &= \text{tr}(\mathbf{D}\mathbf{R}) \hat{\alpha} \mathbb{E} [e^{iuI}] = \text{tr}(\mathbf{D}\mathbf{T}) \hat{\alpha} \mathbb{E} [e^{iuI}] + \mathcal{O}(n^{-1}) \\ &= n\delta\hat{\delta} \mathbb{E} [e^{iuI}] + \mathcal{O}(n^{-1}). \end{aligned} \quad (48)$$

In order to deal with  $\chi_2$ , we apply the results of Proposition 4 related to  $\Psi(\mathbf{Y})$  in the particular case where  $\mathbf{A} = \mathbf{R}$  and  $\mathbf{B} = \tilde{\mathbf{D}}\tilde{\mathbf{R}}$ . In this case,  $\chi_2$  writes  $\chi_2 = iut \mathbb{E} (\Psi(\mathbf{Y})e^{iuI})$ , and Cauchy–Schwarz inequality yields

$$\begin{aligned} |\mathbb{E} (\Psi e^{iuI}) - \mathbb{E} (e^{iuI}) \mathbb{E} (\Psi)| &= |\mathbb{E} [e^{iuI} \hat{\Psi}]| \\ &\leq \sqrt{\mathbb{E} \left[ |\hat{\Psi}(\mathbf{Y})|^2 \right]} = \mathcal{O}(n^{-1}). \end{aligned}$$

Therefore

$$\mathbb{E} (\Psi e^{iuI}) = \mathbb{E} (e^{iuI}) \mathbb{E} (\Psi) + \mathcal{O}(n^{-1}).$$

We now use the approximation for  $\mathbb{E}\Psi(\mathbf{Y})$  given in Proposition 4. By Theorem 3, we can replace  $\tilde{\mathbf{R}}$  (resp.,  $\mathbf{R}$ ) by  $\tilde{\mathbf{T}}$  (resp.,  $\hat{\mathbf{T}}$ ) in the obtained expression. We, therefore, obtain (49) shown at the bottom of the page.

The term  $\chi_3$  can be handled similarly. We apply the results of Proposition 4 related to  $\hat{\Phi}(\mathbf{Y})$  in the particular case where  $\mathbf{A} = \mathbf{R}$  and  $\mathbf{B} = \tilde{\mathbf{D}}\tilde{\mathbf{R}}$ . In this case,  $\chi_3$  writes

$\chi_3 = -tn \mathbb{E} (\hat{\beta} \hat{\Phi}(\mathbf{Y}) e^{iuI})$ , and Cauchy–Schwarz inequality yields

$$\begin{aligned} |\mathbb{E} (\hat{\beta} \hat{\Phi} e^{iuI}) - \mathbb{E} (\hat{\beta} e^{iuI}) \mathbb{E} (\hat{\Phi})| &= |\mathbb{E} [\hat{\beta} e^{iuI} \hat{\Phi}]| \\ &\leq \sqrt{\mathbb{E} [\hat{\beta}^2]} \sqrt{\mathbb{E} [\hat{\Phi}^2]} = \mathcal{O}(n^{-2}). \end{aligned}$$

We therefore obtain

$$\begin{aligned} &\mathbb{E} \left[ \hat{\beta} \text{tr} \left( \mathbf{R}\mathbf{H} \frac{\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\mathbf{Y}^*}{n} \right) e^{iuI} \right] \\ &= \mathbb{E} \left[ \hat{\beta} e^{iuI} \right] \text{tr} (\tilde{\mathbf{D}}^2 \tilde{\mathbf{T}}\tilde{\mathbf{R}}) \frac{1}{n} \text{tr} (\mathbf{D}\mathbf{T}\mathbf{R}) + \mathcal{O}(n^{-1}) \\ &\stackrel{(a)}{=} \mathbb{E} \left[ \hat{\beta} e^{iuI} \right] \tilde{\gamma} \text{tr} (\mathbf{D}\mathbf{T}^2) + \mathcal{O}(n^{-1}) \end{aligned} \quad (50)$$

where (a) follows from Theorem 3. It remains to deal with the term  $\mathbb{E} [\hat{\beta} e^{iuI}]$ . To this end, we shall rely on (46) and develop the term  $\mathbb{E} [H_{pp}e^{iuI}]$ . The Resolvent identity yields

$$\mathbb{E} \left[ [\mathbf{H}\mathbf{Y}\mathbf{Y}^*]_{pp} e^{iuI} \right] = \frac{n}{t} \mathbb{E} [e^{iuI}] - \frac{n}{t} \mathbb{E} [H_{pp}e^{iuI}].$$

Plugging this equality into (46) and using  $r_p = (1 + t\hat{\alpha}d_p)^{-1}$ , after some computations, we obtain (51) shown at the bottom of the page, where (a) follows from Theorem 3, Proposition 4, and Proposition 5. We therefore obtain

$$\begin{aligned} \mathbb{E} [\hat{\beta} e^{iuI}] &= -\frac{1}{n} \frac{iut^2}{(1 - t^2\gamma\tilde{\gamma})^2} \\ &\quad \times \left( \tilde{\gamma} \frac{1}{n} \text{tr} (\mathbf{D}^3 \mathbf{T}^3) - t\gamma^2 \frac{1}{n} \text{tr} (\tilde{\mathbf{D}}^3 \tilde{\mathbf{T}}^3) \right) \varphi + \mathcal{O} \left( \frac{1}{n^2} \right). \end{aligned}$$

$$\begin{aligned} \mathbb{E} (\Psi(\mathbf{Y})e^{iuI}) &= \mathbb{E} \Psi(\mathbf{Y}) \mathbb{E} [e^{iuI}] + \mathcal{O}(n^{-1}) \\ &= \frac{1}{1 - t^2\gamma\tilde{\gamma}} \left( \tilde{\gamma} \frac{1}{n} \text{tr} (\mathbf{D}^2 \mathbf{T}^3) - t\gamma \frac{1}{n} \text{tr} (\tilde{\mathbf{D}}^3 \tilde{\mathbf{T}}^3) \right) \frac{1}{n} \text{tr} (\mathbf{D}\mathbf{T}^2) \mathbb{E} [e^{iuI}] + \mathcal{O}(n^{-1}). \end{aligned} \quad (49)$$

$$\begin{aligned} \mathbb{E} [\hat{\beta} e^{iuI}] &= t^2 \mathbb{E} \left[ \hat{\beta} e^{iuI} \frac{1}{n} \text{tr} \left( \mathbf{R}\mathbf{D}\mathbf{H} \frac{\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\mathbf{Y}^*}{n} \right) \right] \\ &\quad - \frac{iut^2}{n} \mathbb{E} \left[ \frac{1}{n} \text{tr} \left( \mathbf{R}\mathbf{D}\mathbf{H}\mathbf{D}\mathbf{H} \frac{\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\mathbf{Y}^*}{n} \right) e^{iuI} \right] + \frac{1}{n} \text{tr} (\mathbf{D}(\mathbf{R} - \mathbb{E}[\mathbf{H}])) \mathbb{E} [e^{iuI}] \\ &\stackrel{(a)}{=} t^2\gamma\tilde{\gamma} \mathbb{E} [\hat{\beta} e^{iuI}] - \frac{1}{n} \frac{iut^2}{(1 - t^2\gamma\tilde{\gamma})} \left( \tilde{\gamma} \frac{1}{n} \text{tr} (\mathbf{D}^3 \mathbf{T}^3) - t\gamma^2 \frac{1}{n} \text{tr} (\tilde{\mathbf{D}}^3 \tilde{\mathbf{T}}^3) \right) \varphi + \mathcal{O}(n^{-2}) \end{aligned} \quad (51)$$

Plugging (51) into (50), and the result together with (48) and (49) into (47), and getting back to (43) and (42), we obtain

$$\frac{\partial \psi_n(u, t)}{\partial t} = -u^2 \eta_n(t) \psi_n(u, t) + \mathcal{O}(n^{-1})$$

where

$$\eta_n(t) = \frac{1}{1 - t^2 \gamma \tilde{\gamma}} \left( -\frac{t^2 \gamma \frac{1}{n} \text{tr}(\tilde{\mathbf{D}}^3 \tilde{\mathbf{T}}^3) \frac{1}{n} \text{tr}(\mathbf{D} \mathbf{T}^2)}{1 - t^2 \gamma \tilde{\gamma}} + t \tilde{\gamma} \frac{1}{n} \text{tr}(\mathbf{D}^2 \mathbf{T}^3) + \frac{t^3 \tilde{\gamma}^2 \frac{1}{n} \text{tr}(\mathbf{D}^3 \mathbf{T}^3) \frac{1}{n} \text{tr}(\mathbf{D} \mathbf{T}^2)}{1 - t^2 \gamma \tilde{\gamma}} \right). \quad (52)$$

Equation (40) is established, and the first step of the proof is completed.

### B. Identification of the Variance

In order to finish the proof, it remains to prove that

$$\eta_n(t) = \frac{1}{2} \frac{d\sigma_n^2(t)}{dt} \text{ where } \sigma_n^2(t) = -\log(1 - t\gamma_n(t)\tilde{\gamma}_n(t)). \quad (53)$$

To this end, we first begin by computing the derivatives of  $\gamma_n(t)$  and  $\tilde{\gamma}_n(t)$ . We shall prove that

$$\frac{d\tilde{\gamma}}{dt} = -2 \frac{\frac{1}{n} \text{tr}(\tilde{\mathbf{D}}^3 \tilde{\mathbf{T}}^3) \frac{1}{n} \text{tr}(\mathbf{D} \mathbf{T}^2)}{1 - t^2 \gamma \tilde{\gamma}}$$

and

$$\frac{d\gamma}{dt} = -2 \frac{\frac{1}{n} \text{tr}(\mathbf{D}^3 \mathbf{T}^3) \frac{1}{n} \text{tr}(\tilde{\mathbf{D}} \tilde{\mathbf{T}}^2)}{1 - t^2 \gamma \tilde{\gamma}}. \quad (54)$$

We only derive  $\frac{d\tilde{\gamma}}{dt}$ , the computations being similar in the other case. We first expand the expression of  $\tilde{\gamma}$ , and obtain

$$\begin{aligned} \frac{d\tilde{\gamma}}{dt} &= \frac{1}{n} \sum_{j=1}^n \tilde{d}_j^2 \frac{d\tilde{T}_{jj}}{dt} = \frac{1}{n} \sum_{j=1}^n \tilde{d}_j^2 \frac{d}{dt} \left( \frac{1}{1 + t\delta(t)\tilde{d}_j} \right)^2 \\ &= -2 \frac{d}{dt} (t\delta(t)) \frac{1}{n} \text{tr}(\tilde{\mathbf{D}}^3 \tilde{\mathbf{T}}^3). \end{aligned} \quad (55)$$

Let us now compute  $\delta'(t)$

$$\delta'(t) = \frac{1}{n} \sum_{i=1}^N d_i \left( \frac{1}{1 + t\tilde{\delta}(t)d_i} \right)' = -\gamma\tilde{\delta}(t) - \gamma t \tilde{\delta}'(t). \quad (56)$$

A similar computation yields  $\tilde{\delta}'(t) = -\tilde{\gamma}\delta(t) - \tilde{\gamma}t\delta'(t)$ . Combining both equations yields

$$\delta' = \frac{t\gamma\tilde{\gamma}\delta - \gamma\tilde{\delta}}{1 - t^2\gamma\tilde{\gamma}}.$$

We now plug this into (55) and obtain

$$\frac{d\tilde{\gamma}}{dt} = -2 \frac{\frac{1}{n} \text{tr}(\tilde{\mathbf{D}}^3 \tilde{\mathbf{T}}^3) (\delta - t\gamma\tilde{\delta})}{1 - t^2\gamma\tilde{\gamma}}. \quad (57)$$

Recall now that the mere definition of  $\mathbf{T}$ ,  $\tilde{\mathbf{T}}$ ,  $\delta$ , and  $\tilde{\delta}$  yields

$$\begin{cases} t\delta\tilde{\mathbf{D}}\tilde{\mathbf{T}} = \mathbf{I} - \tilde{\mathbf{T}} \\ t\tilde{\delta}\mathbf{D}\mathbf{T} = \mathbf{I} - \mathbf{T} \end{cases}. \quad (58)$$

Using (58), we obtain

$$\begin{aligned} n^{-1} \text{tr}(\mathbf{D} \mathbf{T}^2) &= n^{-1} \text{tr}(\mathbf{D} \mathbf{T} (\mathbf{I} - t\tilde{\delta}\mathbf{D}\mathbf{T})) \\ &= \delta - t\tilde{\delta}\gamma \end{aligned} \quad (59)$$

$$\begin{aligned} n^{-1} \text{tr}(\tilde{\mathbf{D}} \tilde{\mathbf{T}}^2) &= n^{-1} \text{tr}(\tilde{\mathbf{D}} \tilde{\mathbf{T}} (\mathbf{I} - t\delta\tilde{\mathbf{D}}\tilde{\mathbf{T}})) \\ &= \tilde{\delta} - t\delta\tilde{\gamma}. \end{aligned} \quad (60)$$

It remains to plug (59) in (57) to conclude the proof of (54).

We are now in position to prove (53). The main idea in the following computations is to express (52) as a symmetric quantity with respect to  $\delta$  and  $\mathbf{T}$  on the one hand and  $\tilde{\delta}$  and  $\tilde{\mathbf{T}}$ , on the other hand. To this end, we split  $\eta_n(t)$  in (52) as  $\eta_n(t) = \frac{1}{1 - t^2\gamma\tilde{\gamma}} (\eta^{(1)} + \eta^{(2)} + \eta^{(3)})$ . We first work on  $\eta^{(3)}$

$$\begin{aligned} \eta^{(3)} &\stackrel{(a)}{=} \frac{t^3 \delta \tilde{\gamma}^2 \frac{1}{n} \text{tr}(\mathbf{D}^3 \mathbf{T}^3)}{1 - t^2 \gamma \tilde{\gamma}} - \frac{t^4 \tilde{\delta} \tilde{\gamma}^2 \gamma \frac{1}{n} \text{tr}(\mathbf{D}^3 \mathbf{T}^3)}{1 - t^2 \gamma \tilde{\gamma}} \\ &\stackrel{(b)}{=} \frac{-t^2 \tilde{\gamma} \frac{1}{n} \text{tr}(\mathbf{D}^3 \mathbf{T}^3) \frac{1}{n} \text{tr}(\tilde{\mathbf{D}} \tilde{\mathbf{T}}^2)}{1 - t^2 \gamma \tilde{\gamma}} + t^2 \tilde{\gamma} \tilde{\delta} \frac{1}{n} \text{tr}(\mathbf{D}^3 \mathbf{T}^3). \end{aligned}$$

where (a) follows from (59), and (b) from (60). We now look at  $\eta^{(2)}$  as shown in the first equation at the bottom of the page, where the last equality follows (58) again. We therefore have the second equation at the bottom of the page, where (a) follows from (54). This concludes the identification of the variance.

### C. Integration of the Differential (40)

Let us introduce  $K_n(u, \rho) = \psi_n(u, \rho) e^{\frac{u^2}{2} \sigma_n^2(\rho)}$ . Due to (40),  $K_n(u, \rho)$  readily satisfies the following differential equation:

$$\frac{\partial K_n(u, t)}{\partial t} = \varepsilon_n(u, t) e^{\frac{u^2}{2} \sigma_n^2(t)}. \quad (61)$$

$$\eta^{(2)} + t^2 \tilde{\gamma} \tilde{\delta} \frac{1}{n} \text{tr}(\mathbf{D}^3 \mathbf{T}^3) = t \tilde{\gamma} \left( \frac{1}{n} \text{tr}(\mathbf{D}^2 \mathbf{T}^3 + \frac{1}{n} \text{tr}(\mathbf{D}^2 \mathbf{T}^2 (t\tilde{\delta}\mathbf{D}\mathbf{T})) \right) = t\gamma\tilde{\gamma}$$

$$\begin{aligned} \eta_n(t) &= \frac{1}{1 - t^2 \gamma \tilde{\gamma}} \left( -\frac{t^2 \gamma \frac{1}{n} \text{tr}(\tilde{\mathbf{D}}^3 \tilde{\mathbf{T}}^3) \frac{1}{n} \text{tr}(\mathbf{D} \mathbf{T}^2)}{1 - t^2 \gamma \tilde{\gamma}} - \frac{t^2 \tilde{\gamma} \frac{1}{n} \text{tr}(\mathbf{D}^3 \mathbf{T}^3) \frac{1}{n} \text{tr}(\tilde{\mathbf{D}} \tilde{\mathbf{T}}^2)}{1 - t^2 \gamma \tilde{\gamma}} + t\gamma\tilde{\gamma} \right) \\ &\stackrel{(a)}{=} \frac{1}{2} \frac{t^2 \gamma \tilde{\gamma}' + t^2 \gamma' \tilde{\gamma} + 2t\gamma\tilde{\gamma}}{1 - t^2 \gamma \tilde{\gamma}} \\ &= -\frac{1}{2} \frac{d}{dt} \log(1 - t^2 \gamma \tilde{\gamma}) \end{aligned}$$

As in Section IV-C, one can easily prove that  $|\varepsilon_n(t)| \leq \frac{K}{n}$  for every  $t \in [0, \rho]$ . As  $K_n(u, 0) = 1$ , we get

$$K_n(u, \rho) = 1 + \int_0^\rho \varepsilon_n(u, t) e^{\frac{u}{2} \sigma_n^2(t)} dt.$$

Due to Proposition 5,  $\sigma_n^2(t)$  is bounded from above uniformly in  $n$  and  $t \in [0, \rho]$ . This fact, together with  $|\varepsilon_n(t)| \leq \frac{K}{n}$  implies that

$$K_n(u, \rho) = 1 + \mathcal{O}\left(\frac{1}{n}\right).$$

This in turn yields

$$\begin{aligned} \Psi_n(u, \rho) &= (1 + \mathcal{O}(n^{-1})) e^{-\frac{u}{2} \sigma_n^2(\rho)} \\ &= e^{-\frac{u}{2} \sigma_n^2(\rho)} + \mathcal{O}(n^{-1}) \end{aligned}$$

where the last equality follows from the fact that  $\sigma_n^2(\rho)$  is uniformly bounded by  $n$  by Proposition 2.

## APPENDIX

### A. Proof of Proposition 1

Let us first establish the existence and uniqueness of the solution of (2). To this end, we plug the expression of  $\tilde{\delta}$  in (2). The system of two equations reduces to the single equation  $\delta = f(t, \delta)$  where  $f(t, \delta)$  is defined by

$$\begin{aligned} f(t, \delta) &= \frac{1}{n} \text{tr} \left( \mathbf{D}_n \left( \mathbf{I}_N + \frac{t}{n} \text{tr} \right. \right. \\ &\quad \left. \left. \times \left( \tilde{\mathbf{D}}_n \left( \mathbf{I}_n + t \delta \tilde{\mathbf{D}}_n \right)^{-1} \right) \mathbf{D}_n \right)^{-1} \right) \end{aligned} \quad (62)$$

which is itself equivalent to  $g(\delta, t) = 1$  where we get the equation at the bottom of the page. The function  $\delta \mapsto g(t, \delta)$  is continuous, decreasing and satisfies  $g(t, 0) = +\infty$  and  $g(t, +\infty) = 0$ . Therefore, the equation  $g(t, \delta) = 1$  has a unique solution  $\delta(t) > 0$ .

The integral representation (3) of  $\delta$  and  $\tilde{\delta}$  is related to the Stieltjes representation of a class of analytic functions. We refer to [9, Sec. 3.2] where a more general result is proven and skip the details. Be aware however that  $\delta$  in this paper and  $\delta$  in [9] slightly differ (but are related by  $\delta_{\text{IEEE}}(z) = z^{-1} \delta_{\text{AAP}}(-z^{-1})$ ), so do the  $\tilde{\delta}$ 's.

### B. Proof of Proposition 2

In order to prove Proposition 2, it is sufficient to first prove that  $1 - t^2 \gamma \tilde{\gamma}$  is bounded away from zero and then to prove that the same quantity is strictly lower than 1, uniformly in  $n$ . We shall proceed into four steps.

1) *A priori* estimates for  $\delta$ ,  $\tilde{\delta}$ ,  $\gamma$  and  $\tilde{\gamma}$ : The mere definition of  $\delta$  and  $\tilde{\delta}$  yields

$$\delta = \frac{1}{n} \sum_{i=1}^N \frac{d_i}{1 + t d_i \delta} \leq \frac{N d_{\max}}{n}$$

and

$$\tilde{\delta} = \frac{1}{n} \sum_{j=1}^n \frac{\tilde{d}_j}{1 + t \tilde{d}_j \tilde{\delta}} \leq \tilde{d}_{\max}. \quad (63)$$

Using these upper estimates, one gets the following lower estimates:

$$\delta \geq \frac{\frac{1}{n} \text{tr} \mathbf{D}_n}{1 + t d_{\max} \tilde{d}_{\max}} \quad \text{and} \quad \tilde{\delta} \geq \frac{\frac{1}{n} \text{tr} \tilde{\mathbf{D}}_n}{1 + \frac{n}{N} t d_{\max} \tilde{d}_{\max}}. \quad (64)$$

One can notice that due to Assumption (A2), these lower bounds are uniformly bounded in  $n$  away from zero. Finally a straightforward application of Jensen's inequality yields:

$$\delta^2 = \left( \frac{1}{n} \sum_{i=1}^N d_i T_{ii} \right)^2 \leq \frac{N \gamma}{n} \text{ i.e. } \frac{n}{N} \delta^2 \leq \gamma$$

$$\text{Similarly } \tilde{\delta}^2 \leq \tilde{\gamma}. \quad (65)$$

2) An estimate over  $\frac{d\tilde{\delta}}{dt}$ . The following equalities are straightforward (see, for instance, (56)):

$$\delta'(t) = -\gamma \tilde{\delta}(t) - \gamma t \tilde{\delta}'(t)$$

and

$$\tilde{\delta}'(t) = -\tilde{\gamma} \delta(t) - \tilde{\gamma} t \delta'(t). \quad (66)$$

In particular,  $|\tilde{\delta}'(0)| = \tilde{\gamma}(0) \delta(0) \leq N n^{-1} \tilde{d}_{\max}^2 d_{\max}$  which is eventually bounded. Recall that  $\tilde{\delta}$  admits the following representation:

$$\tilde{\delta}(t) = \int_0^\infty \frac{\tilde{\mu}(d\lambda)}{1 + t\lambda}$$

where  $\tilde{\mu}$  is a nonnegative measure satisfying  $\tilde{\mu}(\mathbb{R}^+) = \frac{1}{n} \text{tr} \tilde{\mathbf{D}}_n$ . In particular, one obtains

$$0 < -\tilde{\delta}'(t) = \int_0^\infty \frac{\lambda \tilde{\mu}(d\lambda)}{(1 + t\lambda)^2} \leq -\tilde{\delta}'(0) \leq N n^{-1} \tilde{d}_{\max}^2 d_{\max}. \quad (67)$$

3) The quantity  $1 - t^2 \gamma \tilde{\gamma}$  is bounded away from zero, uniformly in  $n$  and for  $t \in [0, \rho]$ . Eliminating  $\delta'$  between the two equations in (66) yields

$$\begin{aligned} \frac{d\tilde{\delta}}{dt} (1 - t^2 \gamma \tilde{\gamma}) &= \tilde{\gamma} (t \tilde{\delta} \gamma - \delta) = \frac{\tilde{\gamma}}{n} \text{tr} \mathbf{D}_n \mathbf{T}_n \left( t \tilde{\mathbf{D}}_n \mathbf{T}_n - \mathbf{I}_N \right) \\ &= -\frac{\tilde{\gamma}}{n} \text{tr} \mathbf{D}_n \mathbf{T}_n^2 \end{aligned}$$

$$g(t, \delta) = \frac{f(t, \delta)}{\delta} = \frac{1}{n} \text{tr} \left( \mathbf{D}_n \left( \delta \mathbf{I}_N + \frac{t}{n} \text{tr} \left( \delta \tilde{\mathbf{D}}_n \left( \mathbf{I}_n + t \delta \tilde{\mathbf{D}}_n \right)^{-1} \right) \mathbf{D}_n \right)^{-1} \right).$$

where the last equality follows from the identity  $\mathbf{T}_n = (\mathbf{I}_N + t\tilde{\delta}\mathbf{D}_n)^{-1}$  which yields  $(t\tilde{\delta}\mathbf{D}_n\mathbf{T}_n - \mathbf{I}_N) = -\mathbf{T}_n$ . Otherwise stated

$$1 - t^2\gamma\tilde{\gamma} = \frac{\tilde{\gamma}\text{tr}\mathbf{D}_n\mathbf{T}_n^2}{n(-\tilde{\delta}'(t))}.$$

This immediately implies that  $1 - t^2\gamma\tilde{\gamma}$  is positive. In order to check that it is bounded away from zero uniformly in  $n$ , notice first that  $n^{-1}\text{tr}\mathbf{D}_n\mathbf{T}_n^2 \geq d_{\max}^{-1}\gamma$ . Collecting now the previous estimates (65) and (67), we obtain

$$1 - t^2\gamma\tilde{\gamma} \geq \frac{n^2}{N^2} \frac{\delta^2\tilde{\delta}^2}{d_{\max}^2\tilde{d}_{\max}^2}.$$

Using (64) and Assumption (A1), we obtain that  $1 - t^2\gamma\tilde{\gamma}$  is bounded away from zero, uniformly in  $n$  and for  $t \in [0, \rho]$ .

- 4) The quantity  $1 - t^2\gamma\tilde{\gamma}$  is strictly bounded above from 1, uniformly in  $n$ : The inequalities (65) together with (64) yield

$$\sup_n (1 - t^2\gamma\tilde{\gamma}) \leq \sup_n \left(1 - t^2 \frac{n}{N} \delta^2 \tilde{\delta}^2\right) < 1.$$

This completes the proof of Proposition 2.

### C. Proof of Theorem 3

We first give a sketch of the proof to emphasize the main ideas over the technical aspects of the proof.

- 1) We first prove that the asymptotic behavior of  $n^{-1}\text{tr}(\mathbf{A}_n(\mathbf{R}_n - \mathbf{T}_n))$  is directly related to the behavior of  $\alpha(t) - \delta(t)$ . Similarly,  $n^{-1}\text{tr}\tilde{\mathbf{A}}_n(\tilde{\mathbf{R}}_n - \tilde{\mathbf{T}}_n)$  is related to  $\tilde{\alpha}(t) - \tilde{\delta}(t)$ .
- 2) We extend the definition of  $\alpha$  from  $t \in \mathbb{R}^+$  to  $z \in \mathbb{C} \setminus \mathbb{R}_-$  and establish an integral representation

$$\alpha(t) = \int_{\mathbb{R}^+} \frac{\nu(d\lambda)}{1 + \lambda t}.$$

As a consequence of the integral representations for  $\delta$ ,  $\tilde{\delta}$  and  $\alpha$ , we prove that  $\delta$ ,  $\tilde{\delta}$  and  $\alpha$  are bounded analytic functions on every compact subset of  $\mathbb{C} \setminus \mathbb{R}_-$ .

- 3) As a consequence of this detour in the complex plane, we prove the following weaker result. For every uniformly bounded diagonal matrix  $\mathbf{A}_n$ , the following holds true:

$$\begin{cases} n^{-1}\text{tr}(\mathbf{A}_n\mathbf{R}_n) = n^{-1}\text{tr}(\mathbf{A}_n\mathbf{T}_n) + o(1) \\ n^{-1}\text{tr}(\mathbf{A}_n\tilde{\mathbf{R}}_n) = n^{-1}\text{tr}(\tilde{\mathbf{A}}_n\tilde{\mathbf{T}}_n) + o(1). \end{cases}$$

- 4) We then refine the previous result in order to get the sharper rate of convergence  $\mathcal{O}(n^{-2})$  instead of  $o(1)$ .

The theorem will then be proved.

1) *The Asymptotic Behavior of  $n^{-1}\text{tr}(\mathbf{A}_n(\mathbf{R}_n - \mathbf{T}_n))$  and its Relation With  $\alpha(t) - \delta(t)$ :* The standard matrix identity

$$\mathbf{R}_n - \mathbf{T}_n = \mathbf{R}_n(\mathbf{T}_n^{-1} - \mathbf{R}_n^{-1})\mathbf{T}_n$$

immediately yields

$$n^{-1}\text{tr}(\mathbf{A}_n(\mathbf{R}_n - \mathbf{T}_n)) = t(\tilde{\delta}(t) - \tilde{\alpha}(t)) \frac{1}{n} \text{tr}(\mathbf{A}_n\mathbf{R}_n\mathbf{D}_n\mathbf{T}_n)$$

and

$$\begin{aligned} n^{-1}\text{tr}(\tilde{\mathbf{D}}_n(\tilde{\mathbf{T}}_n - \tilde{\mathbf{R}}_n)) &= \tilde{\delta}(t) - \tilde{\alpha}(t) \\ &= t(\alpha(t) - \delta(t)) \frac{1}{n} \text{tr}(\tilde{\mathbf{D}}_n\tilde{\mathbf{R}}_n\tilde{\mathbf{D}}_n\tilde{\mathbf{T}}_n). \end{aligned}$$

Therefore

$$\begin{aligned} n^{-1}\text{tr}(\mathbf{A}_n(\mathbf{R}_n - \mathbf{T}_n)) &= t^2(\alpha(t) - \delta(t)) \\ &\times \frac{1}{n} \text{tr}(\tilde{\mathbf{D}}_n\tilde{\mathbf{R}}_n\tilde{\mathbf{D}}_n\tilde{\mathbf{T}}_n) \frac{1}{n} \text{tr}(\mathbf{A}_n\mathbf{R}_n\mathbf{D}_n\mathbf{T}_n). \end{aligned} \quad (68)$$

2) *An Integral Representation for  $\alpha$ , and Bounds Over  $\alpha$ ,  $\delta$  and  $\tilde{\delta}$ :* Recall that  $\alpha(t) = \mathbb{E}[n^{-1}\text{tr}(\mathbf{D}_n(\mathbf{I}_N + t\mathbf{Y}_n\mathbf{Y}_n^*)^{-1})]$ . This function readily extends from  $t \in \mathbb{R}^+$  to  $z \in \mathbb{C} \setminus \mathbb{R}^-$ . Moreover, the following representation holds true:

$$\alpha(z) = \int_0^{+\infty} \frac{\nu(d\lambda)}{1 + \lambda z} \quad (69)$$

where  $\nu$  is a uniquely defined positive measure on  $\mathbb{R}^+$  such that  $\nu(\mathbb{R}^+) = \frac{1}{n}\text{tr}\mathbf{D}_n$ . To prove this, we introduce the eigenvalue/eigenvector decomposition of matrix  $n^{-1}\mathbf{Y}_n\mathbf{Y}_n^* = \sum_{i=1}^N \lambda_i \mathbf{u}_i \mathbf{u}_i^*$  where  $(\lambda_i, 1 \leq i \leq N)$  and  $(\mathbf{u}_i, 1 \leq i \leq N)$  represent its eigenvalues and eigenvectors, respectively. The random variable  $\beta(z) = \frac{1}{n}\text{tr}\mathbf{D}_n(\mathbf{I} + z\frac{\mathbf{Y}_n\mathbf{Y}_n^*}{n})^{-1}$  can be written as

$$\beta(z) = \frac{1}{n} \sum_{i=1}^N \frac{\mathbf{u}_i^* \mathbf{D}_n \mathbf{u}_i}{\lambda_i - z} = \int_0^{+\infty} \frac{\omega(d\lambda)}{1 + \lambda z}$$

where  $\omega$  is the nonnegative random measure defined by

$$\omega = \frac{1}{n} \sum_{i=1}^N \mathbf{u}_i^* \mathbf{D}_n \mathbf{u}_i \delta(\lambda - \lambda_i).$$

Consider now the measure  $\nu$  defined by  $\nu = \mathbb{E}[\omega]$ , that is  $\nu(B) = \mathbb{E}[\omega(B)]$  for every Borel set  $B \subset \mathbb{R}^+$ . It is clear that  $\alpha(z) = \mathbb{E}[\beta(z)]$  is given by (69), and that  $\nu(\mathbb{R}^+) = \mathbb{E}[\omega(\mathbb{R}^+)]$  is given by

$$\nu(\mathbb{R}^+) = \mathbb{E} \left[ \frac{1}{n} \sum_{i=1}^N \mathbf{u}_i^* \mathbf{D}_n \mathbf{u}_i \right] = \mathbb{E} \left[ \frac{1}{n} \text{tr} \mathbf{D}_n (\sum_i \mathbf{u}_i \mathbf{u}_i^*) \right].$$

As  $\sum_i \mathbf{u}_i \mathbf{u}_i^* = \mathbf{I}_N$ ,  $\nu(\mathbb{R}^+) = \frac{1}{n}\text{tr}\mathbf{D}_n$  as expected and representation (69) implies that  $\alpha(z)$  is analytic over  $\mathbb{C} \setminus \mathbb{R}^-$ .

Let  $\text{dist}(w, \mathbb{R}^+)$  stand for the distance from element  $w \in \mathbb{C}$  to  $\mathbb{R}^+$ . Then the following holds true for every  $z \in \mathbb{C} \setminus \mathbb{R}^-$ :

$$\begin{aligned} |\alpha(z)| &\leq \frac{1}{n} \text{tr}(\mathbf{D}_n) \frac{1}{|z| \text{dist}(-\frac{1}{z}, \mathbb{R}^+)} \\ &\leq \frac{N}{n} d_{\max} \frac{1}{|z| \text{dist}(-\frac{1}{z}, \mathbb{R}^+)}. \end{aligned} \quad (70)$$

Similarly, (3) yields that

$$|\delta(z)| \leq \frac{N d_{\max}}{n|z|} \frac{1}{\text{dist}(-\frac{1}{z}, \mathbb{R}^+)}. \quad (71)$$

A similar result holds for  $\tilde{\delta}(z)$ . These upper bounds imply in particular that  $\alpha(z)$ ,  $\delta(z)$  and  $\tilde{\delta}(z)$  are uniformly bounded on each compact subset of  $\mathbb{C} \setminus \mathbb{R}_-$ .

3) *A Weaker Result as a Consequence of Montel's Theorem:* We first establish that for every diagonal matrix  $\mathbf{A}_n$  uniformly bounded

$$\begin{cases} n^{-1}\text{tr}(\mathbf{A}_n\mathbf{R}_n) = n^{-1}\text{tr}(\mathbf{A}_n\mathbf{T}_n) + o(1) \\ n^{-1}\text{tr}(\tilde{\mathbf{A}}_n\tilde{\mathbf{R}}_n) = n^{-1}\text{tr}(\tilde{\mathbf{A}}_n\tilde{\mathbf{T}}_n) + o(1) \end{cases} \quad (72)$$

We take (68) as a starting point. Matrices  $\mathbf{R}_n$ ,  $\tilde{\mathbf{R}}_n$ ,  $\mathbf{T}_n$ , and  $\tilde{\mathbf{T}}_n$  have their spectral norms bounded by one for  $t \in \mathbb{R}^+$  and matrices  $\mathbf{A}_n$ ,  $\mathbf{D}_n$ , and  $\tilde{\mathbf{D}}_n$  are also uniformly bounded by assumption. Therefore, the terms  $n^{-1}\text{tr}(\tilde{\mathbf{A}}_n\tilde{\mathbf{R}}_n\tilde{\mathbf{D}}_n\tilde{\mathbf{T}}_n)$  and  $n^{-1}\text{tr}(\mathbf{A}_n\mathbf{R}_n\mathbf{D}_n\mathbf{T}_n)$  are also bounded. In order to prove (72), it is sufficient to prove that  $\alpha(t) - \delta(t) = o(1)$ . To this end, we make use of Proposition 5 and write  $\alpha(t) - \delta(t)$  as

$$\alpha(t) - \delta(t) = \frac{1}{n}\text{tr}(\mathbf{D}_n(\mathbf{R}_n - \mathbf{T}_n)) + \varepsilon_n(t),$$

where  $\varepsilon_n(t) = \mathcal{O}(n^{-2})$ . Using relation (68) for  $\mathbf{A}_n = \mathbf{D}_n$ , we immediately get that

$$\alpha(t) - \delta(t) = (\alpha(t) - \delta(t))t^2 \frac{1}{n}\text{tr}(\tilde{\mathbf{D}}_n\tilde{\mathbf{R}}_n\tilde{\mathbf{D}}_n\tilde{\mathbf{T}}_n) \times \frac{1}{n}\text{tr}(\mathbf{D}_n\mathbf{R}_n\mathbf{D}_n\mathbf{T}_n) + \varepsilon_n(t). \quad (73)$$

As  $\sup_n (\|\mathbf{R}_n\|, \|\tilde{\mathbf{R}}_n\|, \|\mathbf{T}_n\|, \|\tilde{\mathbf{T}}_n\|) \leq 1$ , we have

$$\begin{aligned} \frac{1}{n}\text{tr}(\tilde{\mathbf{D}}_n\tilde{\mathbf{R}}_n\tilde{\mathbf{D}}_n\tilde{\mathbf{T}}_n) \frac{1}{n}\text{tr}(\mathbf{D}_n\mathbf{R}_n\mathbf{D}_n\mathbf{T}_n) &\leq \frac{N}{n}d_{\max}^2\tilde{d}_{\max}^2 \\ &\leq 2cd_{\max}^2\tilde{d}_{\max}^2 \end{aligned}$$

as soon as  $\frac{N}{n} \leq 2\ell^+$ , where  $\ell^+$  is defined in (1). Therefore, if  $t < t_0 := (2d_{\max}\tilde{d}_{\max}\sqrt{\ell^+})^{-1}$ , then

$$t^2 \frac{1}{n}\text{tr}(\tilde{\mathbf{D}}_n\tilde{\mathbf{R}}_n\tilde{\mathbf{D}}_n\tilde{\mathbf{T}}_n) \frac{1}{n}\text{tr}(\mathbf{D}_n\mathbf{R}_n\mathbf{D}_n\mathbf{T}_n) < \frac{1}{2}$$

for  $n$  large enough. Equation (73) thus implies that

$$\begin{aligned} |\alpha_n(t) - \delta_n(t)| &< 2|\varepsilon_n(t)|, \text{ i.e.,} \\ \alpha(t) - \delta(t) &= \mathcal{O}(n^{-2}), \text{ for } t < t_0. \end{aligned} \quad (74)$$

This in particular implies that  $\alpha_n(t) - \delta_n(t) = o(1)$  for  $t < t_0$ ; however, it remains to establish this convergence for  $t \geq t_0$ . To this end, observe that  $\alpha_n(z) - \delta_n(z)$  is analytic in  $\mathbb{C} \setminus \mathbb{R}_-$  and bounded on each compact subset of  $\mathbb{C} \setminus \mathbb{R}_-$ . Montel's theorem asserts that the sequence of functions  $\alpha_n(z) - \delta_n(z)$  is compact and therefore that there exists a converging subsequence which converges toward an analytic function. Since this limiting function is zero on  $[0, t_0[$  by (74), it must be zero everywhere due to the analyticity. Therefore from every subsequence, one can extract a subsequence that converges toward zero. Necessarily,  $\alpha_n(z) - \delta_n(z)$  converges to zero for every  $z \in \mathbb{C} \setminus \mathbb{R}^-$  and in particular for  $t \geq 0$ . This establishes (72).

Even if the convergence rate of  $\alpha_n(t) - \delta_n(t)$  is  $\mathcal{O}(n^{-2})$  for  $t < t_0$ , Montel's theorem does not imply that the convergence rate of  $\alpha_n(z) - \delta_n(z)$  remains  $\mathcal{O}(n^{-2})$  elsewhere. Therefore, there remains some work to be done in order to prove that  $\alpha_n(t) - \delta_n(t) = \mathcal{O}(n^{-2})$  for each  $t > 0$ .

4) *End of the Proof:* We take (73) as a starting point. Equation (72) imply that for each  $t \geq 0$

$$\begin{cases} n^{-1}\text{tr}(\mathbf{D}_n\mathbf{R}_n(t)\mathbf{D}_n\mathbf{T}_n(t)) - \gamma(t) = o(1) \\ n^{-1}\text{tr}(\tilde{\mathbf{D}}_n\tilde{\mathbf{R}}_n(t)\tilde{\mathbf{D}}_n\tilde{\mathbf{T}}_n(t)) - \tilde{\gamma}(t) = o(1). \end{cases} \quad (75)$$

where  $\gamma_n = n^{-1}\text{tr}\mathbf{D}_n^2\mathbf{T}_n^2$  and  $\tilde{\gamma}_n = n^{-1}\text{tr}\tilde{\mathbf{D}}_n^2\tilde{\mathbf{T}}_n^2$ . Thanks to Proposition 6, (75) implies that

$$\inf_n \left( 1 - t^2 \frac{1}{n}\text{tr}(\mathbf{D}_n\mathbf{R}_n(t)\mathbf{D}_n\mathbf{T}_n(t)) \times \frac{1}{n}\text{tr}(\tilde{\mathbf{D}}_n\tilde{\mathbf{R}}_n(t)\tilde{\mathbf{D}}_n\tilde{\mathbf{T}}_n(t)) \right) > 0.$$

Equation (73) thus clearly implies that  $\alpha(t) - \delta(t)$  is of the same order of magnitude as  $\varepsilon_n(t)$ , i.e., that  $\alpha(t) - \delta(t) = \mathcal{O}(n^{-2})$ . Theorem 3 is proved.

#### D. Proof of Proposition 4 -(1)—Variance Controls

Consider first  $\Phi(\mathbf{Y}) = \frac{1}{n}\text{tr}(\mathbf{A}\mathbf{H}\frac{\mathbf{Y}\mathbf{B}\mathbf{Y}^*}{n})$ . We use Poincaré-Nash inequality (20) to control the variance of  $\Phi$ . It writes

$$\begin{aligned} \mathbb{E}[\hat{\Phi}(\mathbf{Y})^2] &\leq \sum_{i=1}^N \sum_{j=1}^n d_i \tilde{d}_j \mathbb{E} \left[ \left| \frac{\partial \Phi}{\partial Y_{i,j}} \right|^2 \right] \\ &\quad + \sum_{i=1}^N \sum_{j=1}^n d_i \tilde{d}_j \mathbb{E} \left[ \left| \frac{\partial \Phi}{\partial \bar{Y}_{i,j}} \right|^2 \right]. \end{aligned} \quad (76)$$

We have  $\Phi(\mathbf{Y}) = (1/n^2) \sum_{p,r=1}^N \sum_{q=1}^n a_p b_q H_{pr} Y_{rq} \bar{Y}_{pq}$ . From the differentiation formula (14), we have

$$\begin{aligned} \frac{\partial}{\partial Y_{ij}} (H_{pr} Y_{rq} \bar{Y}_{pq}) &= -\frac{t}{n} H_{pi} [y_j^* \mathbf{H}]_r Y_{rq} \bar{Y}_{pq} \\ &\quad + H_{pr} \bar{Y}_{pq} \delta(r-i) \delta(q-j). \end{aligned}$$

Therefore, after a straightforward computation, we obtain  $\partial\Phi/\partial Y_{ij} = \phi_{ij}^{(1)} + \phi_{ij}^{(2)}$  with

$$\phi_{ij}^{(1)} = -\frac{t}{n^3} [y_j^* \mathbf{H} \mathbf{Y} \mathbf{B} \mathbf{Y}^* \mathbf{A} \mathbf{H}]_i$$

and

$$\phi_{ij}^{(2)} = \frac{1}{n^2} b_j [y_j^* \mathbf{A} \mathbf{H}]_i.$$

The first term of the right-hand side of inequality (76) can be treated as follows:

$$\begin{aligned} &\sum_{i=1}^N \sum_{j=1}^n d_i \tilde{d}_j \mathbb{E} \left[ \left| \frac{\partial \Phi}{\partial Y_{i,j}} \right|^2 \right] \\ &\leq 2 \sum_{i=1}^N \sum_{j=1}^n d_i \tilde{d}_j \left( \mathbb{E} \left[ \left| \phi_{ij}^{(1)} \right|^2 \right] + \mathbb{E} \left[ \left| \phi_{ij}^{(2)} \right|^2 \right] \right) \\ &= \frac{2t^2}{n^6} \mathbb{E} \left[ \text{tr} \left( \mathbf{H} \mathbf{Y} \mathbf{B} \mathbf{Y}^* \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H} \mathbf{A} \mathbf{Y} \mathbf{B} \mathbf{Y}^* \mathbf{H} \mathbf{Y} \tilde{\mathbf{D}} \mathbf{Y}^* \right) \right] \\ &\quad + \frac{2}{n^4} \mathbb{E} \left[ \text{tr} \left( \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H} \mathbf{A} \mathbf{Y} \mathbf{B}^2 \tilde{\mathbf{D}} \mathbf{Y}^* \right) \right]. \end{aligned} \quad (77)$$

Let  $A = \sup \|\mathbf{A}\|$ . Using inequalities (8), (9) and (11) and Cauchy–Schwarz inequality, we have (78) shown at the bottom of the page, where the last inequality is due to (12). Turning to the second term of the right-hand side of (77), we have (79) shown at the bottom of the page. The second term of the right-hand side of Inequality (76) is treated similarly. This proves that  $\text{var}(\Phi) = \mathcal{O}(n^{-2})$ .

Consider now  $\Psi(\mathbf{Y}) = \frac{1}{n} \text{tr} \left( \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H} \frac{\mathbf{Y} \mathbf{B} \mathbf{Y}^*}{n} \right)$ . The proof being quite similar to the previous one, we just give its main steps. By (20) we have  $\mathbb{E}[\hat{\Psi}(\mathbf{Y})^2] \leq \sum_{i=1}^N \sum_{j=1}^n d_i \tilde{d}_j (\mathbb{E}[|\partial \Psi / \partial Y_{i,j}|^2] + \mathbb{E}[|\partial \Psi / \partial \bar{Y}_{i,j}|^2])$ . A computation similar to above yields  $\partial \Psi / \partial Y_{ij} = \psi_{ij}^{(1)} + \psi_{ij}^{(2)} + \psi_{ij}^{(3)}$  where

$$\begin{aligned} \psi_{ij}^{(1)} &= -\frac{t}{n^3} [\mathbf{y}_j^* \mathbf{H} \mathbf{D} \mathbf{H} \mathbf{Y} \mathbf{B} \mathbf{Y}^* \mathbf{A} \mathbf{H}]_i \\ \psi_{ij}^{(2)} &= -\frac{t}{n^3} [\mathbf{y}_j^* \mathbf{H} \mathbf{Y} \mathbf{B} \mathbf{Y}^* \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H}]_i \\ \psi_{ij}^{(3)} &= \frac{1}{n^2} b_j [\mathbf{y}_j^* \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H}]_i. \end{aligned}$$

We have the third equation at the bottom of the page. The first two terms of the right-hand side can be bounded by a series of inequalities similar to inequalities (78). The third term can be bounded as in (79). This ends the proofs of the variance controls in Proposition 4.

#### E. Proof of Proposition 4-(2)—Approximation Rules

Consider first  $\Phi(Y) = \frac{1}{n} \text{tr} \left( \mathbf{A} \mathbf{H} \frac{\mathbf{Y} \mathbf{B} \mathbf{Y}^*}{n} \right)$ . We write

$$\Phi(\mathbf{Y}) = n^{-2} \sum_{p,i=1}^N \sum_{j=1}^n a_p b_j \mathbb{E} [Y_{ij} H_{pi} \bar{Y}_{pj}]$$

and apply the Integration by parts formula (19) to the summand. Using identity (15), we have the fourth equation at the bottom of the page. By taking the sum over the index  $i$ , we obtain

$$\mathbb{E} \left[ [\mathbf{H} \mathbf{y}_j]_p \bar{Y}_{pj} \right] = -t \tilde{d}_j \mathbb{E} \left[ \beta [\mathbf{H} \mathbf{y}_j]_p \bar{Y}_{pj} \right] + d_p \tilde{d}_j \mathbb{E} [H_{pp}].$$

$$\begin{aligned} & \frac{2t^2}{n^6} \mathbb{E} \left[ \text{tr} \left( \mathbf{H} \mathbf{Y} \mathbf{B} \mathbf{Y}^* \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H} \mathbf{A} \mathbf{Y} \mathbf{B} \mathbf{Y}^* \mathbf{H} \mathbf{Y} \tilde{\mathbf{D}} \mathbf{Y}^* \right) \right] \\ & \leq \frac{2t^2}{n^6} \mathbb{E} \left[ \sqrt{\text{tr} \left( (\mathbf{H} \mathbf{Y} \mathbf{B} \mathbf{Y}^* \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H} \mathbf{A} \mathbf{Y} \mathbf{B} \mathbf{Y}^* \mathbf{H})^2 \right)} \sqrt{\text{tr} \left( (\mathbf{Y} \tilde{\mathbf{D}} \mathbf{Y}^*)^2 \right)} \right] \\ & \leq \frac{2t^2}{n^6} \mathbb{E} \left[ \|\mathbf{H}\|^4 \|\mathbf{A}\|^2 \|\mathbf{D}\| \sqrt{\text{tr} \left( (\mathbf{Y} \mathbf{B} \mathbf{Y}^*)^4 \right)} \sqrt{\text{tr} \left( (\mathbf{Y} \tilde{\mathbf{D}} \mathbf{Y}^*)^2 \right)} \right] \\ & \leq \frac{2d_{\max} A^2 t^2}{n^2} \sqrt{\frac{1}{n} \mathbb{E} \left[ \text{tr} \left( \left( \frac{\mathbf{Y} \mathbf{B} \mathbf{Y}^*}{n} \right)^4 \right) \right]} \sqrt{\frac{1}{n} \mathbb{E} \left[ \text{tr} \left( \left( \frac{\mathbf{Y} \tilde{\mathbf{D}} \mathbf{Y}^*}{n} \right)^2 \right) \right]} \\ & < \frac{K}{n^2} \end{aligned} \tag{78}$$

$$\begin{aligned} \frac{2}{n^4} \mathbb{E} \left[ \text{tr} \left( \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H} \mathbf{A} \mathbf{Y} \mathbf{B}^2 \tilde{\mathbf{D}} \mathbf{Y}^* \right) \right] & \leq \frac{2A^2 d_{\max}}{n^2} \mathbb{E} \left[ \frac{1}{n} \text{tr} \left( \frac{1}{n} \mathbf{Y} \mathbf{B}^2 \tilde{\mathbf{D}} \mathbf{Y}^* \right) \right] \\ & < \frac{K}{n^2}. \end{aligned} \tag{79}$$

$$\begin{aligned} \sum_{i=1}^N \sum_{j=1}^n d_i \tilde{d}_j \mathbb{E} \left[ \left| \frac{\partial \Psi}{\partial Y_{i,j}} \right|^2 \right] & \leq 3 \sum_{i=1}^N \sum_{j=1}^n d_i \tilde{d}_j \left( \mathbb{E} \left[ |\psi_{ij}^{(1)}|^2 \right] + \mathbb{E} \left[ |\psi_{ij}^{(2)}|^2 \right] + \mathbb{E} \left[ |\psi_{ij}^{(3)}|^2 \right] \right) \\ & = + \frac{3t^2}{n^6} \mathbb{E} \left[ \text{tr} \left( \mathbf{H} \mathbf{D} \mathbf{H} \mathbf{Y} \mathbf{B} \mathbf{Y}^* \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H} \mathbf{A} \mathbf{Y} \mathbf{B} \mathbf{Y}^* \mathbf{H} \mathbf{D} \mathbf{H} \mathbf{Y} \tilde{\mathbf{D}} \mathbf{Y}^* \right) \right] \\ & \quad + \frac{3t^2}{n^6} \mathbb{E} \left[ \text{tr} \left( \mathbf{H} \mathbf{Y} \mathbf{B} \mathbf{Y}^* \mathbf{A} (\mathbf{H} \mathbf{D})^3 \mathbf{H} \mathbf{A} \mathbf{Y} \mathbf{B} \mathbf{Y}^* \mathbf{H} \mathbf{Y} \tilde{\mathbf{D}} \mathbf{Y}^* \right) \right] \\ & \quad + \frac{3}{n^4} \mathbb{E} \left[ \text{tr} \left( \mathbf{A} (\mathbf{H} \mathbf{D})^3 \mathbf{H} \mathbf{A} \mathbf{Y} \mathbf{B}^2 \tilde{\mathbf{D}} \mathbf{Y}^* \right) \right]. \end{aligned}$$

$$\mathbb{E} [Y_{ij} H_{pi} \bar{Y}_{pj}] = d_i \tilde{d}_j \mathbb{E} \left[ \frac{\partial}{\partial Y_{ij}} (H_{pi} \bar{Y}_{pj}) \right] = -\frac{t}{n} d_i \tilde{d}_j \mathbb{E} \left[ [\mathbf{H} \mathbf{y}_j]_p H_{ii} \bar{Y}_{pj} \right] + d_i \tilde{d}_j \delta(i-p) \mathbb{E} [H_{pi}].$$

Writing now  $\beta = \hat{\beta} + \alpha$  and then grouping together the terms with  $\mathbb{E} \left[ \left[ \mathbf{H}\mathbf{y}_j \right]_p \overline{Y_{pj}} \right]$ , we obtain

$$\mathbb{E} \left[ \left[ \mathbf{H}\mathbf{y}_j \right]_p \overline{Y_{pj}} \right] = -t\tilde{d}_j\tilde{r}_j\mathbb{E} \left[ \hat{\beta} \left[ \mathbf{H}\mathbf{y}_j \right]_p \overline{Y_{pj}} \right] + d_p\tilde{d}_j\tilde{r}_j\mathbb{E} [H_{pp}].$$

We now sum over  $j$  and  $p$ , and obtain

$$\mathbb{E} \left[ \frac{1}{n} \text{tr} \left( \mathbf{A}\mathbf{H} \frac{\mathbf{Y}\mathbf{B}\mathbf{Y}^*}{n} \right) \right] = \frac{1}{n} \text{tr} \left( \tilde{\mathbf{D}}\tilde{\mathbf{R}}\tilde{\mathbf{B}} \right) \frac{1}{n} \text{tr} \left( \mathbf{A}\mathbf{D} \mathbb{E} [\mathbf{H}] \right) + \varepsilon$$

with

$$\begin{aligned} \varepsilon &= -t\mathbb{E} \left[ \hat{\beta} \frac{1}{n} \text{tr} \left( \mathbf{A}\mathbf{H} \frac{\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\tilde{\mathbf{B}}\mathbf{Y}^*}{n} \right) \right] \\ &= -t\mathbb{E} \left[ \hat{\beta} \frac{1}{n} \text{tr} \left( \mathbf{A}\mathbf{H} \frac{\widehat{\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\tilde{\mathbf{B}}\mathbf{Y}^*}}{n} \right) \right]. \end{aligned}$$

Applying Cauchy–Schwarz inequality, Proposition 3 and the variance controls in Proposition 4, we get  $|\varepsilon| = \mathcal{O}(n^{-2})$ .

By Theorem 3,  $n^{-1}\text{tr}(\tilde{\mathbf{D}}\tilde{\mathbf{R}}\tilde{\mathbf{B}}) = n^{-1}\text{tr}(\tilde{\mathbf{D}}\tilde{\mathbf{T}}\tilde{\mathbf{B}}) + \mathcal{O}(n^{-2})$ . By Theorem 3 and Proposition 5, we obtain  $n^{-1}\text{tr}(\mathbf{A}\mathbf{D} \mathbb{E} [\mathbf{H}]) = n^{-1}\text{tr}(\mathbf{A}\mathbf{D}\mathbf{T}) + \mathcal{O}(n^{-2})$ . This ends the proof of (23).

Consider now  $\Psi(\mathbf{Y}) = \frac{1}{n}\text{tr}(\mathbf{A}\mathbf{H}\mathbf{D}\mathbf{H} \frac{\mathbf{Y}\mathbf{B}\mathbf{Y}^*}{n})$ . In order to compute  $\mathbb{E}\Psi(\mathbf{Y})$ , we shall need the following intermediate result:

*Lemma 1:* In the setting of Theorem 1, let  $\Upsilon(\mathbf{Y}) = \frac{1}{n}\text{tr}(\mathbf{D}\mathbf{H}\mathbf{D}\mathbf{H})$ . Then

1) The following estimate holds true:

$$\text{var}[\Upsilon(\mathbf{Y})] = \mathcal{O}\left(\frac{1}{n^2}\right);$$

2) moreover

$$\mathbb{E}[\Upsilon(\mathbf{Y})] = \frac{\gamma}{1-t^2\gamma\tilde{\gamma}} + \mathcal{O}\left(\frac{1}{n^2}\right).$$

*Proof:* In order to prove Lemma 1-(1), we use the Resolvent identity (10) and write

$$\mathbf{D}\mathbf{H}\mathbf{D}\mathbf{H} = \mathbf{D}\mathbf{H}\mathbf{D} - tn^{-1}\mathbf{D}\mathbf{H}\mathbf{D}\mathbf{H}\mathbf{Y}\mathbf{Y}^*.$$

Since  $\text{var}(X+Y) \leq 2\text{var}(X) + 2\text{var}(Y)$ , we only need to deal with each term of the right-hand side. By Proposition 3,  $\text{var}(n^{-1}\text{tr} \mathbf{D}\mathbf{H}\mathbf{D}) = \mathcal{O}(n^{-2})$  and by Proposition 4 -(1),  $\text{var}(tn^{-2}\text{tr} \mathbf{D}\mathbf{H}\mathbf{D}\mathbf{H}\mathbf{Y}\mathbf{Y}^*) = \mathcal{O}(n^{-2})$  and the proof of Lemma 1-(1) is completed.

Let us now prove Lemma 1-(2). The Resolvent identity (10) yields

$$\mathbb{E} \left[ \left[ \mathbf{H}\mathbf{D}\mathbf{H} \right]_{pp} \right] = d_p\mathbb{E} [H_{pp}] - t\mathbb{E} \left[ \left[ \mathbf{H}\mathbf{D}\mathbf{H} \frac{\mathbf{Y}\mathbf{Y}^*}{n} \right]_{pp} \right]. \quad (80)$$

We then write

$$\mathbb{E} \left[ \left[ \mathbf{H}\mathbf{D}\mathbf{H} \frac{\mathbf{Y}\mathbf{Y}^*}{n} \right]_{pp} \right] = n^{-1} \sum_{k,i=1}^N \sum_{j=1}^n d_k H_{pk} H_{ki} Y_{ij} \overline{Y_{pj}}$$

and apply the differentiation formula (14) to the summand. After derivations similar to (44)–(45), we obtain

$$\begin{aligned} &\frac{1}{n}\mathbb{E} \left[ \left[ \mathbf{H}\mathbf{D}\mathbf{H}\mathbf{y}_j \right]_p \overline{Y_{pj}} \right] \\ &= -\frac{t}{n}\tilde{d}_j\tilde{r}_j\mathbb{E} \left[ \left[ \mathbf{H}\mathbf{y}_j \right]_p \overline{Y_{pj}} \frac{1}{n} \text{tr}(\mathbf{D}\mathbf{H}\mathbf{D}\mathbf{H}) \right] \\ &\quad - \frac{t}{n}\tilde{d}_j\tilde{r}_j\mathbb{E} \left[ \hat{\beta} \left[ \mathbf{H}\mathbf{D}\mathbf{H}\mathbf{y}_j \right]_p \overline{Y_{pj}} \right] \\ &\quad + \frac{1}{n}d_p\tilde{d}_j\tilde{r}_j\mathbb{E} \left[ \left[ \mathbf{H}\mathbf{D}\mathbf{H} \right]_{pp} \right]. \end{aligned} \quad (81)$$

Taking the sum over  $j$  and combining with (80) yields

$$\begin{aligned} \mathbb{E} \left[ \left[ \mathbf{H}\mathbf{D}\mathbf{H} \right]_{pp} \right] &= t^2 r_p \mathbb{E} \left[ \left[ \mathbf{H} \frac{\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\mathbf{Y}^*}{n} \right]_{pp} \frac{1}{n} \text{tr}(\mathbf{D}\mathbf{H}\mathbf{D}\mathbf{H}) \right] \\ &\quad + t^2 r_p \mathbb{E} \left[ \hat{\beta} \left[ \mathbf{H}\mathbf{D}\mathbf{H} \frac{\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\mathbf{Y}^*}{n} \right]_{pp} \right] \\ &\quad + r_p d_p \mathbb{E} [H_{pp}]. \end{aligned} \quad (82)$$

Taking now the sum over  $p$ , we obtain

$$\begin{aligned} \mathbb{E} \left[ \frac{1}{n} \text{tr}(\mathbf{D}\mathbf{H}\mathbf{D}\mathbf{H}) \right] &= \frac{1}{n} \sum_{p=1}^N d_p \mathbb{E} \left[ \left[ \mathbf{H}\mathbf{D}\mathbf{H} \right]_{pp} \right] \\ &= \chi_1 + \chi_2 + \chi_3 \end{aligned} \quad (83)$$

where

$$\begin{aligned} \chi_1 &= +t^2 \mathbb{E} \left[ \frac{1}{n} \text{tr} \left( \mathbf{D}\mathbf{R}\mathbf{H} \frac{\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\mathbf{Y}^*}{n} \right) \frac{1}{n} \text{tr}(\mathbf{D}\mathbf{H}\mathbf{D}\mathbf{H}) \right] \\ \chi_2 &= t^2 \mathbb{E} \left[ \hat{\beta} \frac{1}{n} \text{tr} \left( \mathbf{D}\mathbf{R}\mathbf{H}\mathbf{D}\mathbf{H} \frac{\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\mathbf{Y}^*}{n} \right) \right] \\ \chi_3 &= \frac{1}{n} \text{tr}(\mathbf{D}^2\mathbf{R} \mathbb{E} [\mathbf{H}]). \end{aligned}$$

Let us first deal with the terms  $\chi_2$  and  $\chi_3$ . Cauchy–Schwarz inequality together with Proposition 3 and Proposition 4-(1) yield  $\chi_2 = \mathcal{O}(n^{-2})$ . Proposition 5 together with Theorem 3 yield  $\chi_3 = \gamma + \mathcal{O}(n^{-2})$ . We now look at  $\chi_1$ . Due to Proposition 4-(1) and to Lemma 1-(1), we have

$$\begin{aligned} \chi_1 &= t^2 \mathbb{E} \left[ \frac{1}{n} \text{tr} \left( \mathbf{D}\mathbf{R}\mathbf{H} \frac{\mathbf{Y}\tilde{\mathbf{D}}\tilde{\mathbf{R}}\mathbf{Y}^*}{n} \right) \right] \\ &\quad \times \mathbb{E} \left[ \frac{1}{n} \text{tr}(\mathbf{D}\mathbf{H}\mathbf{D}\mathbf{H}) \right] + \mathcal{O}\left(\frac{1}{n^2}\right), \\ &\stackrel{(a)}{=} t^2\gamma\tilde{\gamma} \mathbb{E} \left[ \frac{1}{n} \text{tr}(\mathbf{D}\mathbf{H}\mathbf{D}\mathbf{H}) \right] + \mathcal{O}\left(\frac{1}{n^2}\right) \end{aligned}$$



$$\begin{aligned} \mathbb{E} \left[ \frac{1}{n} \text{tr} \left( \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H} \frac{\mathbf{Y} \mathbf{B} \mathbf{Y}^*}{n} \right) \right] &= -t \mathbb{E} \left[ \frac{1}{n} \text{tr} (\mathbf{D} \mathbf{H} \mathbf{D} \mathbf{H}) \frac{1}{n} \text{tr} \left( \mathbf{A} \mathbf{H} \frac{\mathbf{Y} \tilde{\mathbf{D}} \tilde{\mathbf{R}} \mathbf{B} \mathbf{Y}^*}{n} \right) \right] \\ &+ \frac{t^2}{n} \text{tr} (\tilde{\mathbf{D}} \tilde{\mathbf{R}} \mathbf{B}) \mathbb{E} \left[ \frac{1}{n} \text{tr} (\mathbf{D} \mathbf{H} \mathbf{D} \mathbf{H}) \frac{1}{n} \text{tr} \left( \mathbf{A} \mathbf{R} \mathbf{D} \mathbf{H} \frac{\mathbf{Y} \tilde{\mathbf{D}} \tilde{\mathbf{R}} \mathbf{Y}^*}{n} \right) \right] \\ &+ \frac{1}{n} \text{tr} (\tilde{\mathbf{D}} \tilde{\mathbf{R}} \mathbf{B}) \frac{1}{n} \text{tr} (\mathbf{A} \mathbf{D}^2 \mathbf{R} \mathbf{E} \mathbf{H}) + \mathcal{O} \left( \frac{1}{n^2} \right). \end{aligned}$$

$$\begin{aligned} \mathbb{E} \left[ \frac{1}{n} \text{tr} \left( \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H} \frac{\mathbf{Y} \mathbf{B} \mathbf{Y}^*}{n} \right) \right] &= -t \mathbb{E} \left[ \frac{1}{n} \text{tr} (\mathbf{D} \mathbf{H} \mathbf{D} \mathbf{H}) \right] \mathbb{E} \left[ \frac{1}{n} \text{tr} \left( \mathbf{A} \mathbf{H} \frac{\mathbf{Y} \tilde{\mathbf{D}} \tilde{\mathbf{R}} \mathbf{B} \mathbf{Y}^*}{n} \right) \right] \\ &+ t^2 \frac{1}{n} \text{tr} (\tilde{\mathbf{D}} \tilde{\mathbf{R}} \mathbf{B}) \mathbb{E} \left[ \frac{1}{n} \text{tr} (\mathbf{D} \mathbf{H} \mathbf{D} \mathbf{H}) \right] \mathbb{E} \left[ \frac{1}{n} \text{tr} \left( \mathbf{A} \mathbf{R} \mathbf{D} \mathbf{H} \frac{\mathbf{Y} \tilde{\mathbf{D}} \tilde{\mathbf{R}} \mathbf{Y}^*}{n} \right) \right] \\ &+ \frac{1}{n} \text{tr} (\tilde{\mathbf{D}} \tilde{\mathbf{R}} \mathbf{B}) \frac{1}{n} \text{tr} (\mathbf{A} \mathbf{D}^2 \mathbf{R} \mathbf{E} \mathbf{H}) + \mathcal{O} \left( \frac{1}{n^2} \right). \end{aligned}$$

where (a) follows from (23) in Proposition 4. It remains to plug the values obtained for  $\chi_1$ ,  $\chi_2$  and  $\chi_3$  into (83) to obtain

$$(1 - t^2 \gamma \tilde{\gamma}) \mathbb{E} \left[ \frac{1}{n} \text{tr} (\mathbf{D} \mathbf{H} \mathbf{D} \mathbf{H}) \right] = \gamma + \mathcal{O}(n^{-2}).$$

Recalling Proposition 2, we can divide by  $(1 - t^2 \gamma \tilde{\gamma})$  and obtain the desired result.  $\square$

We can now go back to the computation of  $\mathbb{E} \Psi(\mathbf{Y})$ . Let us give the main steps of the derivation. Expanding  $\mathbb{E} \Psi(\mathbf{Y})$  yields:

$$\begin{aligned} \mathbb{E} \left[ \frac{1}{n} \text{tr} \left( \mathbf{A} \mathbf{H} \mathbf{D} \mathbf{H} \frac{\mathbf{Y} \mathbf{B} \mathbf{Y}^*}{n} \right) \right] &= \frac{1}{n^2} \\ &\times \sum_{p=1}^N \sum_{j=1}^n a_p b_j \mathbb{E} \left[ \left[ \mathbf{H} \mathbf{D} \mathbf{H} \mathbf{y}_j \right]_p \overline{Y_{pj}} \right]. \end{aligned}$$

We replace the summand  $n^{-1} \mathbb{E} \left[ \left[ \mathbf{H} \mathbf{D} \mathbf{H} \mathbf{y}_j \right]_p \overline{Y_{pj}} \right]$  by the expression given by (81). We then replace the term  $\mathbb{E} \left[ \left[ \mathbf{H} \mathbf{D} \mathbf{H} \right]_{pp} \right]$  in (81) by the expression given by (82). We sum over  $p$  and  $j$  and notice afterwards that the terms where  $\hat{\beta}$  is involved are of order  $\mathcal{O}(n^{-2})$ . We therefore end up with the first equation at the top of the page. We first decorrelate by using the variance estimates in Proposition 4-(1) and Lemma 1-(1) and obtain the second equation at the top of the page. It remains to apply Theorem 3, Proposition 4 and Lemma 1-(2) to the terms in the right-hand side of the previous equality to conclude.

#### REFERENCES

[1] I. E. Telatar, "Capacity of multi-antenna Gaussian channels," *European Trans. Telecommun.*, vol. 10, no. 6, pp. 585–595, Nov./Dec. 1999.  
 [2] G. J. Foschini, "Layered space-time architecture for wireless communication in a fading environment when using multiple antennas," *Bell Labs Tech. J.*, vol. 1, no. 2, pp. 41–59, 1996.  
 [3] V. A. Marchenko and L. A. Pastur, "Distribution of eigenvalues for some sets of random matrices," *Math. USSR—Sbornik*, vol. 1, no. 4, pp. 457–483, 1967.

[4] R. R. Müller, "A random matrix model of communication via antenna arrays," *IEEE Trans. Inf. Theory*, vol. 48, no. 9, pp. 2495–2506, Sep. 2002.  
 [5] C.-N. Chuah, D. N. C. Tse, J. M. Kahn, and R. A. Valenzuela, "Capacity scaling in MIMO wireless systems under correlated fading," *IEEE Trans. Inf. Theory*, vol. 48, no. 3, pp. 637–650, Mar. 2002.  
 [6] X. Mestre, J. R. Fonollosa, and A. Pagès-Zamora, "Capacity of MIMO channels: Asymptotic evaluation under correlated fading," *IEEE J. Sel. Areas Commun.*, vol. 21, pp. 829–838, Jun. 2003.  
 [7] A. L. Moustakas, S. H. Simon, and A. M. Sengupta, "MIMO capacity through correlated channels in the presence of correlated interferers and noise: A (Not So) large N analysis," *IEEE Trans. Inf. Theory*, vol. 49, no. 10, pp. 2545–2561, Oct. 2003.  
 [8] A. M. Tulino, A. Lozano, and S. Verdú, "Impact of antenna correlation on the capacity of multi-antenna channels," *IEEE Trans. Inf. Theory*, vol. 51, no. 7, pp. 2491–2509, Jul. 2005.  
 [9] W. Hachem, P. Loubaton, and J. Najim, "Deterministic equivalents for certain functionals of large random matrices," *Ann. Appl. Probab.*, vol. 17, no. 3, pp. 875–930, 2007.  
 [10] W. Hachem, P. Loubaton, and J. Najim, "The empirical eigenvalue distribution of a gram matrix: from independence to stationarity," *Markov Process. Rel. Fields*, vol. 11, no. 4, pp. 629–648, 2005.  
 [11] D.-S. Shiu, G. J. Foschini, M. J. Gans, and J. M. Kahn, "Fading correlation and its effect on the capacity of multielement antenna systems," *IEEE Trans. Commun.*, vol. 48, no. 3, pp. 502–513, Mar. 2000.  
 [12] J. P. Kermaol, L. Schumacher, K. I. Pedersen, P. E. Mogensen, and F. Frederiken, "A stochastic MIMO radio channel model with experimental validation," *IEEE J. Sel. Areas Commun.*, vol. 20, pp. 1211–1225, 2002.  
 [13] W. Hachem, P. Loubaton, and J. Najim, "A CLT for information-theoretic statistics of Gram random matrices with a given variance profile," *Ann. Appl. Probab.*, 2007.  
 [14] G. W. Anderson and O. Zeitouni, "A CLT for a band matrix model," *Prob. Theory Rel. Fields*, vol. 134, no. 2, pp. 283–338, 2006.  
 [15] Z. D. Bai and J. W. Silverstein, "CLT for linear spectral statistics of large-dimensional sample covariance matrices," *Ann. Probab.*, vol. 32, no. 1A, pp. 553–605, 2004.  
 [16] A. B. de Monvel and A. Khorunzhy, "Limit theorems for random matrices," *Markov Process. Rel. Fields*, vol. 4, no. 2, pp. 175–197, 1998.  
 [17] A. M. Khorunzhy and L. A. Pastur, "Limits of infinite interaction radius, dimensionality and the number of components for random operators with off-diagonal randomness," *Commun. Math. Phys.*, vol. 153, no. 3, pp. 605–646, 1993.  
 [18] L. A. Pastur, A. M. Khorunzhi, and V. Y. Vasil'chuk, "On an asymptotic property of the spectrum of the sum of one-dimensional independent random operators," *Dopov. Nats. Akad. Nauk Ukraïni*, no. 2, pp. 27–30, 1995.  
 [19] L. A. Pastur, "A simple approach to the global regime of Gaussian ensembles of random matrices," *Ukrainian Math. J.*, vol. 57, no. 6, pp. 936–966, Jun. 2005.

- [20] L. H. Y. Chen, "An inequality for the multivariate normal distribution," *J. Multivariate Anal.*, vol. 12, no. 2, pp. 306–315, 1982.
- [21] C. Houdré, V. Pérez-Abreu, and D. Surgailis, "Interpolation, correlation identities, and inequalities for infinitely divisible variables," *J. Fourier Anal. Appl.*, vol. 4, no. 6, pp. 651–668, 1998.
- [22] S. Chatterjee and A. Bose, "A new method for bounding rates of convergence of empirical spectral distributions," *J. Theoret. Probab.*, vol. 17, no. 4, pp. 1003–1019, 2004.
- [23] S. Chatterjee, "Fluctuations of eigenvalues and second order Poincaré inequalities," *Probab. Theory Related Fields*, 2007, to be published.
- [24] A. Guionnet and O. Zeitouni, "Concentration of the spectral measure for large matrices," *Electron. Commun. Probab.*, vol. 5, pp. 119–136, 2000.
- [25] J. Dumont, W. Hachem, S. Lasaulce, P. Loubaton, and J. Najim, "On the capacity achieving covariance matrix for rician MIMO channels: An asymptotic approach," *IEEE Trans. Inf. Theory*, 2007, submitted for publication.
- [26] R. Horn and C. Johnson, *Matrix Analysis*. Cambridge, U.K.: Cambridge Univ. Press, 1994.
- [27] J. Glimm and A. Jaffe, *Quantum physics*, second ed. New York: Springer-Verlag, 1987.
- [28] P. Billingsley, *Probability and Measure*, 3rd ed. New York: Wiley, 1995.